


# MAP3K5 & APP Protein Protein Interaction Antibody Pair

Catalog # DI0061 Size 1 Set

## Applications



Representative image of Proximity Ligation Assay of protein-protein interactions between MAP3K5 and APP. HeLa cells were stained with anti-MAP3K5 rabbit purified polyclonal antibody 1:1200 and anti-APP mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

## Specification

<b>Product Description</b>	This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the MAP3K5 protein, and the other against the APP protein for use in <a href="#">in situ Proximity Ligation Assay</a> . See Publication Reference below.
<b>Reactivity</b>	Human
<b>Quality Control Testing</b>	Protein protein interaction immunofluorescence result. Representative image of Proximity Ligation Assay of protein-protein interactions between MAP3K5 and APP. HeLa cells were stained with anti-MAP3K5 rabbit purified polyclonal antibody 1:1200 and anti-APP mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.
<b>Supplied Product</b>	Antibody pair set content: 1. MAP3K5 rabbit purified polyclonal antibody (100 ug) 2. APP mouse monoclonal antibody (40 ug) *Reagents are sufficient for at least 30-50 assays using recommended protocols.
<b>Storage Instruction</b>	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

## Applications

- *In situ* Proximity Ligation Assay (Cell)

## Gene Info — APP

Entrez GeneID	<a href="#">351</a>
Gene Name	APP
Gene Alias	AAA, ABETA, ABPP, AD1, APPI, CTFgamma, CVAP, PN2
Gene Description	amyloid beta (A4) precursor protein
Omim ID	<a href="#">104760 605714</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	This gene encodes a cell surface receptor and transmembrane precursor protein that is cleaved by secretases to form a number of peptides. Some of these peptides are secreted and can bind to the acetyltransferase complex APBB1/TIP60 to promote transcriptional activation, while others form the protein basis of the amyloid plaques found in the brains of patients with Alzheimer disease. Mutations in this gene have been implicated in autosomal dominant Alzheimer disease and cerebral arteriovenous malformations (cerebral amyloid angiopathy). Multiple transcript variants encoding several different isoforms have been found for this gene. [provided by RefSeq]
Other Designations	A4 amyloid protein amyloid beta A4 protein amyloid-beta protein beta-amyloid peptide cerebral vascular amyloid peptide peptidase nexin-II protease nexin-II

## Gene Info — MAP3K5

Entrez GeneID	<a href="#">4217</a>
Gene Name	MAP3K5
Gene Alias	ASK1, MAPKKK5, MEKK5
Gene Description	mitogen-activated protein kinase kinase kinase 5
Omim ID	<a href="#">602448</a>
Gene Ontology	<a href="#">Hyperlink</a>

**Gene Summary**

Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulated kinase (ERK), MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAPKKK or MEKK). MAPKK kinase/MEKK phosphorylates and activates its downstream protein kinase, MAPK kinase/MEK, which in turn activates MAPK. The kinases of these signaling cascades are highly conserved, and homologs exist in yeast, Drosophila, and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. Northern blot analysis shows that MAPKKK5 transcript is abundantly expressed in human heart and pancreas. The MAPKKK5 protein phosphorylates and activates MKK4 (aliases SERK1, MAPKK4) in vitro, and activates c-Jun N-terminal kinase (JNK)/stress-activated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK. [provided by RefSeq]

**Other Designations**

MAP/ERK kinase kinase 5|MAPK/ERK kinase kinase 5|OTTHUMP00000017275|apoptosis signal regulating kinase

**Pathway**

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [MAPK signaling pathway](#)
- [Neurotrophin signaling pathway](#)

**Disease**

- [Alzheimer disease](#)
- [Amyloidosis](#)
- [Asthma](#)
- [Cardiovascular Diseases](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Cerebral Hemorrhage](#)
- [Cerebrovascular Disorders](#)
- [Cognition](#)
- [Cognition Disorders](#)
- [Dementia](#)

- [Diabetes Mellitus](#)
- [Diabetes Mellitus](#)
  
- [Disease Progression](#)
  
- [Disease Susceptibility](#)
  
- [Down Syndrome](#)
  
- [Edema](#)
  
- [Edema](#)
  
- [Genetic Predisposition to Disease](#)
  
- [Genetic Predisposition to Disease](#)
  
- [Headache](#)
  
- [Hypersensitivity](#)
  
- [Inflammation](#)
  
- [Insulin Resistance](#)
  
- [Lymphoma](#)
  
- [Macular Degeneration](#)
  
- [Mental Status Schedule](#)
  
- [Neuropsychological Tests](#)
  
- [Psychiatric Status Rating Scales](#)
  
- [Recurrence](#)
  
- [Tobacco Use Disorder](#)
  
- [Tobacco Use Disorder](#)
  
- [Tourette Syndrome](#)