


# TRAF2 & ERN1 Protein Protein Interaction Antibody Pair

Catalog # DI0168      Size 1 Set

## Applications



Representative image of Proximity Ligation Assay of protein-protein interactions between TRAF2 and ERN1. HeLa cells were stained with anti-TRAF2 rabbit purified polyclonal antibody 1:1200 and anti-ERN1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

## Specification

<b>Product Description</b>	This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the TRAF2 protein, and the other against the ERN1 protein for use in <a href="#">in situ Proximity Ligation Assay</a> . <a href="#">See Publication Reference below</a> .
<b>Reactivity</b>	Human
<b>Quality Control Testing</b>	Protein protein interaction immunofluorescence result. Representative image of Proximity Ligation Assay of protein-protein interactions between TRAF2 and ERN1. HeLa cells were stained with anti-TRAF2 rabbit purified polyclonal antibody 1:1200 and anti-ERN1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.
<b>Supplied Product</b>	Antibody pair set content: 1. TRAF2 rabbit purified polyclonal antibody (100 ug) 2. ERN1 mouse monoclonal antibody (40 ug) *Reagents are sufficient for at least 30-50 assays using recommended protocols.
<b>Storage Instruction</b>	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze-thaw cycle. Reagents should be returned to -20°C storage immediately after use.

## Applications

- *In situ* Proximity Ligation Assay (Cell)

## Gene Info — ERN1

<b>Entrez GeneID</b>	<a href="#">2081</a>
<b>Gene Name</b>	ERN1
<b>Gene Alias</b>	FLJ30999, IRE1, IRE1P, MGC163277, MGC163279
<b>Gene Description</b>	endoplasmic reticulum to nucleus signaling 1
<b>Omim ID</b>	<a href="#">604033</a>
<b>Gene Ontology</b>	<a href="#">Hyperlink</a>
<b>Gene Summary</b>	The protein encoded by this gene is the ER to nucleus signalling 1 protein, a human homologue of the yeast Ire1 gene product. This protein possesses intrinsic kinase activity and an endoribonuclease activity and it is important in altering gene expression as a response to endoplasmic reticulum-based stress signals. [provided by RefSeq]
<b>Other Designations</b>	ER to nucleus signalling 1 endoplasmic reticulum to nucleus signalling 1 inositol-requiring 1 protein kinase/endoribonuclease

## Gene Info — TRAF2

<b>Entrez GeneID</b>	<a href="#">7186</a>
<b>Gene Name</b>	TRAF2
<b>Gene Alias</b>	MGC:45012, TRAP, TRAP3
<b>Gene Description</b>	TNF receptor-associated factor 2
<b>Omim ID</b>	<a href="#">601895</a>
<b>Gene Ontology</b>	<a href="#">Hyperlink</a>

### Gene Summary

The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from members of the TNF receptor superfamily. This protein directly interacts with TNF receptors, and forms a heterodimeric complex with TRAF1. This protein is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this protein and TRAF1 interacts with the inhibitor-of-apoptosis proteins (IAPs), and functions as a mediator of the anti-apoptotic signals from TNF receptors. The interaction of this protein with TRADD, a TNF receptor associated apoptotic signal transducer, ensures the recruitment of IAPs for the direct inhibition of caspase activation. BIRC2/c-IAP1, an apoptosis inhibitor possessing ubiquitin ligase activity, can ubiquitinate and induce the degradation of this protein, and thus potentiate TNF-induced apoptosis. Multiple alternatively spliced transcript variants have been found for this gene, but the biological validity of only one transcript has been determined. [provided by RefSeq]

### Other Designations

OTTHUMP00000022625|OTTHUMP00000064745|tumor necrosis factor type 2 receptor associated protein 3

## Pathway

- [Adipocytokine signaling pathway](#)
- [Apoptosis](#)
- [MAPK signaling pathway](#)
- [Pathways in cancer](#)
- [Small cell lung cancer](#)

## Disease

- [Alzheimer disease](#)
- [Cardiovascular Diseases](#)
- [Cardiovascular Diseases](#)
- [Connective Tissue Diseases](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Diabetes Mellitus](#)
- [Ductus Arteriosus](#)
- [Edema](#)

- [Edema](#)
- [Fetal Diseases](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hematologic Diseases](#)
- [HIV Infections](#)
- [Hodgkin Disease](#)
- [Infant](#)
- [Infection](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Metabolic Syndrome X](#)
- [Multiple Myeloma](#)
- [Musculoskeletal Diseases](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Neoplasms](#)
- [Occupational Diseases](#)
- [Osteoporosis](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Retinopathy of Prematurity](#)
- [Skin Diseases](#)

- [Tobacco Use Disorder](#)
- [Waldenstrom Macroglobulinemia](#)
  
- [Werner syndrome](#)