


AKT1 & BAD Protein Protein Interaction Antibody Pair


Catalog # DI0242 Size 1 Set

Applications




In situ Proximity Ligation Assay (Cell)

Representative image of Proximity Ligation Assay of protein-protein interactions between AKT1 and BAD. HT-29 cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:100 and anti-BAD mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).




In situ Proximity Ligation Assay (Cell)

Representative image of Proximity Ligation Assay of protein-protein interactions between AKT1 and BAD. PC-3 cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:100 and anti-BAD mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).



In situ Proximity Ligation Assay (Cell)

Representative image of Proximity Ligation Assay of protein-protein interactions between AKT1 and BAD. A-549 cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:100 and anti-BAD mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).



Representative image of Proximity Ligation Analysis of protein-protein interactions between AKT1 and BAD. HeLa cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:100 and anti-BAD mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Specification

Product Description	This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the AKT1 protein, and the other against the BAD protein for use in <u>In situ Proximity Ligation Assay</u> . See Publication Reference below.
Reactivity	Human
Quality Control Testing	Protein protein interaction immunofluorescence result. Representative image of Proximity Ligation Analysis of protein-protein interactions between AKT1 and BAD. HeLa cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:100 and anti-BAD mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.
Supplied Product	Antibody pair set content: 1. AKT1 rabbit purified polyclonal antibody (100 ug) 2. BAD mouse monoclonal antibody (40 ug) *Reagents are sufficient for at least 30-50 assays using recommended protocols.
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

- *In situ* Proximity Ligation Assay (Cell)

Representative image of Proximity Ligation Assay of protein-protein interactions between AKT1 and BAD. HT-29 cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:100 and anti-BAD mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

- *In situ* Proximity Ligation Assay (Cell)

Representative image of Proximity Ligation Assay of protein-protein interactions between AKT1 and BAD. PC-3 cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:100 and anti-BAD mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

- *In situ* Proximity Ligation Assay (Cell)

Representative image of Proximity Ligation Assay of protein-protein interactions between AKT1 and BAD. A-549 cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:100 and anti-BAD mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

Gene Info — AKT1

Entrez GenelD	207
Gene Name	AKT1
Gene Alias	AKT, MGC99656, PKB, PKB-ALPHA, PRKBA, RAC, RAC-ALPHA
Gene Description	v-akt murine thymoma viral oncogene homolog 1
Omim ID	164730 181500
Gene Ontology	Hyperlink
Gene Summary	The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidyl inositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq]
Other Designations	RAC-alpha serine/threonine-protein kinase murine thymoma viral (v-akt) oncogene homolog-1 protein kinase B rac protein kinase alpha

Gene Info — BAD

Entrez GenelD	572
Gene Name	BAD
Gene Alias	BBC2, BCL2L8
Gene Description	BCL2-associated agonist of cell death
Omim ID	603167
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. [provided by RefSeq]

Other Designations

BCL-X/BCL-2 binding protein|BCL2-antagonist of cell death protein|BCL2-binding component 6|BCL2-binding protein

Pathway

- [Acute myeloid leukemia](#)
- [Acute myeloid leukemia](#)
- [Adipocytokine signaling pathway](#)
- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Apoptosis](#)
- [B cell receptor signaling pathway](#)
- [Chemokine signaling pathway](#)
- [Chronic myeloid leukemia](#)
- [Chronic myeloid leukemia](#)
- [Colorectal cancer](#)
- [Colorectal cancer](#)
- [Endometrial cancer](#)
- [Endometrial cancer](#)
- [ErbB signaling pathway](#)
- [ErbB signaling pathway](#)
- [Fc epsilon RI signaling pathway](#)
- [Fc gamma R-mediated phagocytosis](#)

- [Focal adhesion](#)
- [Focal adhesion](#)
- [Glioma](#)
- [Insulin signaling pathway](#)
- [Insulin signaling pathway](#)
- [Jak-STAT signaling pathway](#)
- [MAPK signaling pathway](#)
- [Melanoma](#)
- [Melanoma](#)
- [mTOR signaling pathway](#)
- [Neurotrophin signaling pathway](#)
- [Neurotrophin signaling pathway](#)
- [Non-small cell lung cancer](#)
- [Non-small cell lung cancer](#)
- [Pancreatic cancer](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Prostate cancer](#)
- [Renal cell carcinoma](#)
- [Small cell lung cancer](#)
- [T cell receptor signaling pathway](#)
- [Tight junction](#)
- [Toll-like receptor signaling pathway](#)
- [VEGF signaling pathway](#)

- [VEGF signaling pathway](#)

Disease

- [Adenocarcinoma](#)
- [Alzheimer disease](#)
- [Amphetamine-Related Disorders](#)
- [Atherosclerosis](#)
- [Basal Ganglia Diseases](#)
- [Bipolar Disorder](#)
- [Breast Neoplasms](#)
- [Calcinosis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cardiovascular Diseases](#)
- [Cognition](#)
- [Colonic Neoplasms](#)
- [Colorectal Neoplasms](#)
- [Coronary Artery Disease](#)
- [Depressive Disorder](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Dominance](#)
- [Drug Toxicity](#)

- [Dyskinesia](#)
- [Edema](#)
- [Edema](#)
- [Endometrial Neoplasms](#)
- [Endometriosis](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Genetic Predisposition to Disease](#)
- [HIV Infections](#)
- [Leukemia](#)
- [Liver Cirrhosis](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Memory](#)
- [Metabolic Syndrome X](#)
- [Necrosis](#)
- [Neoplasm Metastasis](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Neuropsychological Tests](#)
- [Obesity](#)
- [Osteoporosis](#)
- [Ovarian Failure](#)
- [Ovarian Neoplasms](#)
- [Parkinson disease](#)
- [Parkinson disease](#)

- [Polycystic Ovary Syndrome](#)
- [Precursor T-Cell Lymphoblastic Leukemia-Lymphoma](#)
- [Prostatic Neoplasms](#)
- [Psychiatric Status Rating Scales](#)
- [Psychoses](#)
- [Psychotic Disorders](#)
- [Puberty](#)
- [Pulmonary Disease](#)
- [Rectal Neoplasms](#)
- [Retinal Neoplasms](#)
- [Retinoblastoma](#)
- [Schizophrenia](#)
- [Space Perception](#)
- [Thrombophilia](#)
- [Thyroid Neoplasms](#)
- [Thyroid Neoplasms](#)
- [Tobacco Use Disorder](#)
- [Tuberculosis](#)
- [Urinary Bladder Neoplasms](#)
- [Verbal Learning](#)
- [Werner syndrome](#)