


# MAX & SMAD3 Protein Protein Interaction Antibody Pair

Catalog # DI0382


Size 1 Set

## Applications




### *In situ Proximity Ligation Assay (Cell)*

Representative image of Proximity Ligation Assay of protein-protein interactions between MAX and SMAD3. HT-29 cells were stained with anti-MAX rabbit purified polyclonal antibody 1:100 and anti-SMAD3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).




### *In situ Proximity Ligation Assay (Cell)*

Representative image of Proximity Ligation Assay of protein-protein interactions between MAX and SMAD3. PC-3 cells were stained with anti-MAX rabbit purified polyclonal antibody 1:100 and anti-SMAD3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).




### *In situ Proximity Ligation Assay (Cell)*

Representative image of Proximity Ligation Assay of protein-protein interactions between MAX and SMAD3. A-549 cells were stained with anti-MAX rabbit purified polyclonal antibody 1:100 and anti-SMAD3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).



### *In situ* Proximity Ligation Assay (Cell)

Confocal microscopy image of Proximity Ligation Assay of protein-protein interactions between MAX and SMAD3. PC-3 cells were stained with anti-MAX rabbit purified polyclonal antibody 1:100 and anti-SMAD3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).



Representative image of Proximity Ligation Assay of protein-protein interactions between MAX and SMAD3. HeLa cells were stained with anti-MAX rabbit purified polyclonal antibody 1:1200 and anti-SMAD3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

## Specification

<b>Product Description</b>	This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the MAX protein, and the other against the SMAD3 protein for use in <a href="#">in situ Proximity Ligation Assay</a> . <a href="#">See Publication Reference below</a> .
<b>Reactivity</b>	Human
<b>Quality Control Testing</b>	Protein protein interaction immunofluorescence result. Representative image of Proximity Ligation Assay of protein-protein interactions between MAX and SMAD3. HeLa cells were stained with anti-MAX rabbit purified polyclonal antibody 1:1200 and anti-SMAD3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.
<b>Supplied Product</b>	Antibody pair set content: 1. MAX rabbit purified polyclonal antibody (100 ug) 2. SMAD3 mouse monoclonal antibody (40 ug) *Reagents are sufficient for at least 30-50 assays using recommended protocols.
<b>Storage Instruction</b>	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

## Applications

- *In situ* Proximity Ligation Assay (Cell)

Representative image of Proximity Ligation Assay of protein-protein interactions between MAX and SMAD3. HT-29 cells were stained with anti-MAX rabbit purified polyclonal antibody 1:100 and anti-SMAD3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

- *In situ* Proximity Ligation Assay (Cell)

Representative image of Proximity Ligation Assay of protein-protein interactions between MAX and SMAD3. PC-3 cells were stained with anti-MAX rabbit purified polyclonal antibody 1:100 and anti-SMAD3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

- *In situ* Proximity Ligation Assay (Cell)

Representative image of Proximity Ligation Assay of protein-protein interactions between MAX and SMAD3. A-549 cells were stained with anti-MAX rabbit purified polyclonal antibody 1:100 and anti-SMAD3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

- *In situ* Proximity Ligation Assay (Cell)

Confocal microscopy image of Proximity Ligation Assay of protein-protein interactions between MAX and SMAD3. PC-3 cells were stained with anti-MAX rabbit purified polyclonal antibody 1:100 and anti-SMAD3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

## Gene Info — SMAD3

Entrez GeneID	<a href="#">4088</a>
Gene Name	SMAD3
Gene Alias	DKFZp586N0721, DKFZp686J10186, HSPC193, HsT17436, JV15-2, MADH3, MGC60396
Gene Description	SMAD family member 3
Omim ID	<a href="#">603109</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein functions as a transcriptional modulator activated by transforming growth factor-beta and is thought to play a role in the regulation of carcinogenesis. [provided by RefSeq]
Other Designations	MAD, mothers against decapentaplegic homolog 3 SMA- and MAD-related protein 3 SMAD, mothers against DPP homolog 3 mad homolog JV15-2 mad protein homolog mothers against decapentaplegic homolog 3

## Gene Info — MAX

Entrez GeneID	<a href="#">4149</a>
Gene Name	MAX
Gene Alias	MGC10775, MGC11225, MGC18164, MGC34679, MGC36767, bHLHd4, bHLHd5, bHLHd6, bHLHd7, bHLHd8, orf1
Gene Description	MYC associated factor X
Omim ID	<a href="#">154950</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	The protein encoded by this gene is a member of the basic helix-loop-helix leucine zipper (bHLHZ ) family of transcription factors. It is able to form homodimers and heterodimers with other family members, which include Mad, Mxi1 and Myc. Myc is an oncoprotein implicated in cell proliferation , differentiation and apoptosis. The homodimers and heterodimers compete for a common DNA target site (the E box) and rearrangement among these dimer forms provides a complex system of transcriptional regulation. Multiple alternatively spliced transcript variants have been described for this gene but the full-length nature for some of them is unknown. [provided by RefSeq]
Other Designations	MAX protein helix-loop-helix zipper protein myc-associated factor X

## Pathway

- [Adherens junction](#)
- [Cell cycle](#)
- [Chronic myeloid leukemia](#)
- [Colorectal cancer](#)
- [MAPK signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Pathways in cancer](#)
- [Small cell lung cancer](#)
- [TGF-beta signaling pathway](#)

- [Wnt signaling pathway](#)

## Disease

- [Alzheimer disease](#)
- [Anemia](#)
- [Asthma](#)
- [Bacteremia](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Coronary Artery Disease](#)
- [Crohn Disease](#)
- [Diabetes Mellitus](#)
- [Diabetic Nephropathies](#)
- [Genetic Predisposition to Disease](#)
- [Graft vs Host Disease](#)
- [Head and Neck Neoplasms](#)
- [Hypersensitivity](#)
- [Hypertension](#)
- [Keloid](#)
- [Kidney Failure](#)
- [Liver Cirrhosis](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Obesity](#)

- [Occupational Diseases](#)
- [Osteoarthritis](#)
- [Osteoporosis](#)
- [Ovarian cancer](#)
- [Ovarian Failure](#)
- [Ovarian Neoplasms](#)
- [Pancreatic cancer](#)
- [Pancreatic Neoplasms](#)
- [Polycystic Ovary Syndrome](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Puberty](#)
- [Pulmonary Disease](#)
- [Thrombophilia](#)
- [Tobacco Use Disorder](#)