


# HCK & CRKL Protein Protein Interaction Antibody Pair

Catalog # DI0390 Size 1 Set

# **Applications**



Representative image of Proximity Ligation Assay of protein-protein interactions between HCK and CRKL. Mahlavu cells were stained with anti-HCK rabbit purified polyclonal antibody 1:1200 and anti-CRKL mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Specification	
Product Description	This protein protein interaction antibody pair set comes with two antibodies to detect the protein-prot ein interaction, one against the HCK protein, and the other against the CRKL protein for use in <u>in situ</u> Proximity Ligation Assay. See Publication Reference below.
Reactivity	Human
Quality Control Testing	Protein protein interaction immunofluorescence result.  Representative image of Proximity Ligation Assay of protein-protein interactions between HCK and CRKL. Mahlavu cells were stained with anti-HCK rabbit purified polyclonal antibody 1:1200 and anti-CRKL mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein int eraction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.
Supplied Product	Antibody pair set content:  1. HCK rabbit purified polyclonal antibody (100 ug)  2. CRKL mouse monoclonal antibody (40 ug)  *Reagents are sufficient for at least 30-50 assays using recommended protocols.
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze tha w cycle. Reagents should be returned to -20°C storage immediately after use.

### **Applications**



• In situ Proximity Ligation Assay (Cell)

Gene Info — CRKL	
Entrez GenelD	1399
Gene Name	CRKL
Gene Alias	-
Gene Description	v-crk sarcoma virus CT10 oncogene homolog (avian)-like
Omim ID	602007
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene encodes a protein kinase containing SH2 and SH3 (src homology) domains which has been shown to activate the RAS and JUN kinase signaling pathways and transform fibroblasts in a RAS-dependent fashion. It is a substrate of the BCR-ABL tyrosine kinase, plays a role in fibrobl ast transformation by BCR-ABL, and may be oncogenic
Other Designations	v-crk avian sarcoma virus CT10 oncogene homolog-like

Gene Info — HCK	
Entrez GenelD	<u>3055</u>
Gene Name	HCK
Gene Alias	JTK9
Gene Description	hemopoietic cell kinase
Omim ID	<u>142370</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a protein-tyrosine kinase that is predominantly expressed in hemopoietic cell types. The encoded protein may help couple the Fc receptor to the activation of the respiratory burst. In addition, it may play a role in neutrophil migration and in the degranulation of neutrophils. Alternate translation initiation site usage, including a non-AUG (CUG) codon, result in the production of two different isoforms, that have different subcellular localization. [provided by RefSeq
Other Designations	tyrosine protein kinase HCK



## **Pathway**

- Chemokine signaling pathway
- Chemokine signaling pathway
- Chronic myeloid leukemia
- ErbB signaling pathway
- Fc gamma R-mediated phagocytosis
- Fc gamma R-mediated phagocytosis
- Focal adhesion
- Insulin signaling pathway
- MAPK signaling pathway
- Neurotrophin signaling pathway
- Pathways in cancer
- Regulation of actin cytoskeleton
- Renal cell carcinoma

#### Disease

- Cardiovascular Diseases
- Diabetes Mellitus
- Edema
- HIV Infections
- Pulmonary Disease