


STAT1 & STAT5A Protein Protein Interaction Antibody Pair

Catalog # DI0474

Size 1 Set

Applications



Representative image of Proximity Ligation Assay of protein-protein interactions between STAT1 and STAT5A. HeLa cells were stained with anti-STAT1 rabbit purified polyclonal antibody 1:1200 and anti-STAT5A mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Specification

Product Description

This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the STAT1 protein, and the other against the STAT5A protein for use in [in situ Proximity Ligation Assay](#). [See Publication Reference below](#).

Reactivity

Human

Quality Control Testing

Protein protein interaction immunofluorescence result.
Representative image of Proximity Ligation Assay of protein-protein interactions between STAT1 and STAT5A. HeLa cells were stained with anti-STAT1 rabbit purified polyclonal antibody 1:1200 and anti-STAT5A mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied Product

Antibody pair set content:
1. STAT1 rabbit purified polyclonal antibody (100 ug)
2. STAT5A mouse monoclonal antibody (40 ug)
*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze-thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

- *In situ* Proximity Ligation Assay (Cell)

Gene Info — STAT1

Entrez GeneID	6772
Gene Name	STAT1
Gene Alias	DKFZp686B04100, ISGF-3, STAT91
Gene Description	signal transducer and activator of transcription 1, 91kDa
Omim ID	209950 600555
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the STAT protein family. In response to cytokine s and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. Two alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq]
Other Designations	OTTHUMP00000165047 signal transducer and activator of transcription 1 signal transducer and activator of transcription-1 transcription factor ISGF-3 components p91/p84

Gene Info — STAT5A

Entrez GeneID	6776
Gene Name	STAT5A
Gene Alias	MGF, STAT5
Gene Description	signal transducer and activator of transcription 5A
Omim ID	601511
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for the tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of this gene in cells. [provided by RefSeq]

Other Designations

-

Pathway

- [Acute myeloid leukemia](#)
- [Chemokine signaling pathway](#)
- [Chronic myeloid leukemia](#)
- [ErbB signaling pathway](#)
- [Jak-STAT signaling pathway](#)
- [Jak-STAT signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Pathways in cancer](#)
- [Toll-like receptor signaling pathway](#)

Disease

- [Arthritis](#)
- [Asthma](#)
- [Asthma](#)
- [Birth Weight](#)
- [Birth Weight](#)
- [Breast cancer](#)

- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Breast Neoplasms](#)
- [Bronchiolitis](#)
- [Bronchiolitis](#)
- [Campylobacter Infections](#)
- [Carcinoma](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chronic Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Glioma](#)
- [Glomerulonephritis](#)
- [Hepatitis B](#)
- [Hepatitis C](#)
- [Hepatitis C](#)
- [Hypersensitivity](#)

- [Infant](#)
- [Infant](#)
- [Leukemia](#)
- [Leukemia](#)
- [Liver Cirrhosis](#)
- [Liver Neoplasms](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Lupus Erythematosus](#)
- [Lupus Erythematosus](#)
- [Lymphoma](#)
- [Meningeal Neoplasms](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Meningioma](#)
- [Multiple Sclerosis](#)
- [Neutropenia](#)
- [Osteoporosis](#)
- [Ovarian Neoplasms](#)
- [Papillomavirus Infections](#)
- [Proteinuria](#)
- [Pulmonary Disease](#)
- [Respiratory Syncytial Virus Infections](#)
- [Respiratory Syncytial Virus Infections](#)
- [Salmonella Infections](#)
- [Thrombocytopenia](#)

- [Thyroid Neoplasms](#)
- [Tobacco Use Disorder](#)
- [Tuberculosis](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Viremia](#)
- [Viremia](#)
- [Werner syndrome](#)