


# FGF5 & FGFR4 Protein Protein Interaction Antibody Pair

Catalog # DI0622 Size 1 Set

## Applications



Representative image of Proximity Ligation Assay of protein-protein interactions between FGF5 and FGFR4. HeLa cells were stained with anti-FGF5 rabbit purified polyclonal antibody 1:1200 and anti-FGFR4 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

## Specification

<b>Product Description</b>	This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the FGF5 protein, and the other against the FGFR4 protein for use in <a href="#">in situ Proximity Ligation Assay</a> . <a href="#">See Publication Reference below</a> .
<b>Reactivity</b>	Human
<b>Quality Control Testing</b>	Protein protein interaction immunofluorescence result. Representative image of Proximity Ligation Assay of protein-protein interactions between FGF5 and FGFR4. HeLa cells were stained with anti-FGF5 rabbit purified polyclonal antibody 1:1200 and anti-FGFR4 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.
<b>Supplied Product</b>	Antibody pair set content: 1. FGF5 rabbit purified polyclonal antibody (100 ug) 2. FGFR4 mouse monoclonal antibody (40 ug) *Reagents are sufficient for at least 30-50 assays using recommended protocols.
<b>Storage Instruction</b>	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

## Applications

- *In situ* Proximity Ligation Assay (Cell)

## Gene Info — FGF5

Entrez GenelD	<a href="#">2250</a>
Gene Name	FGF5
Gene Alias	HBGF-5, Smag-82
Gene Description	fibroblast growth factor 5
Omim ID	<a href="#">165190</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This gene was identified as an oncogene, which confers transforming potential when transfected into mammalian cells. Targeted disruption of the homolog of this gene in mouse resulted in the phenotype of abnormally long hair, which suggested a function as an inhibitor of hair elongation. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq]
Other Designations	heparin-binding growth factor 5

## Gene Info — FGFR4

Entrez GenelD	<a href="#">2264</a>
Gene Name	FGFR4
Gene Alias	CD334, JTK2, MGC20292, TKF
Gene Description	fibroblast growth factor receptor 4
Omim ID	<a href="#">134935</a>
Gene Ontology	<a href="#">Hyperlink</a>

## Gene Summary

The protein encoded by this gene is a member of the fibroblast growth factor receptor family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. The genomic organization of this gene, compared to members 1-3, encompasses 18 exons rather than 19 or 20. Although alternative splicing has been observed, there is no evidence that the C-terminal half of the IgIII domain of this protein varies between three alternate forms, as indicated for members 1-3. This particular family member preferentially binds acidic fibroblast growth factor and, although its specific function is unknown, it is overexpressed in gynecological tumor samples, suggesting a role in breast and ovarian tumorigenesis. [provided by RefSeq]

## Other Designations

OTTHUMP00000161430|hydroxaryl-protein kinase|protein-tyrosine kinase|tyrosine kinase related to fibroblast growth factor receptor|tyrosyl|protein kinase

## Pathway

- [Endocytosis](#)
- [MAPK signaling pathway](#)
- [MAPK signaling pathway](#)
- [Melanoma](#)
- [Pathways in cancer](#)
- [Regulation of actin cytoskeleton](#)
- [Regulation of actin cytoskeleton](#)

## Disease

- [ACTH-Secreting Pituitary Adenoma](#)
- [Adenocarcinoma](#)
- [Bone Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)

- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Cleft Palate](#)
- [Colon cancer](#)
- [Colorectal Neoplasms](#)
- [Disease Progression](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Head and Neck Neoplasms](#)
- [Hypersensitivity](#)
- [Hypertension](#)
- [Irritable Bowel Syndrome](#)
- [Kidney Failure](#)
- [Liver Neoplasms](#)
- [Lung carcinoma](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Malignant melanoma](#)
- [Melanoma](#)
- [Mouth Neoplasms](#)
- [Neoplasm Metastasis](#)

- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Pituitary ACTH Hypersecretion](#)
- [Pituitary Neoplasms](#)
- [Prostate cancer](#)
- [Prostatic Hyperplasia](#)
- [Prostatic Neoplasms](#)
- [Recurrence](#)
- [Sarcoma](#)
- [Skin Neoplasms](#)
- [Soft Tissue Neoplasms](#)
- [Thyroid Neoplasms](#)
- [Urinary Bladder Neoplasms](#)