


ACTL6A (Human) Recombinant Protein (Q01)

Catalog # H00000086-Q01

Size 25 ug, 10 ug

Applications



Specification

Product Description	Human ACTL6A partial ORF (NP_817126, 1 a.a. - 80 a.a.) recombinant protein with GST-tag at N-terminal.
Sequence	MVVERDDGSTLMEIDGDKGKQGGPTYIDTNALRVPRENMEAISPLKNGMVEDWDSFQILDHTY KMHVKSEASLHPVLM
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	34.54
Interspecies Antigen Sequence	Mouse (96); Rat (95)
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — ACTL6A

Entrez GeneID [86](#)

GeneBank Accession# [NM_177989](#)

Protein Accession# [NP_817126](#)

Gene Name ACTL6A

Gene Alias ACTL6, ARPN-BETA, Arp4, BAF53A, INO80K, MGC5382

Gene Description actin-like 6A

Omim ID [604958](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene encodes a family member of actin-related proteins (ARPs), which share significant amino acid sequence identity to conventional actins. Both actins and ARPs have an actin fold, which includes an ATP-binding cleft, as a common feature. The ARPs are involved in diverse cellular processes, including vesicular transport, spindle orientation, nuclear migration and chromatin remodeling. This gene encodes a 53 kDa subunit protein of the BAF (BRG1/brm-associated factor) complex in mammals, which is functionally related to SWI/SNF complex in *S. cerevisiae* and *Drosophila*; the latter is thought to facilitate transcriptional activation of specific genes by antagonizing chromatin-mediated transcriptional repression. Together with beta-actin, it is required for maximal ATPase activity of BRG1, and for the association of the BAF complex with chromatin/matrix. Three transcript variants that encode two different protein isoforms have been described. [provided by RefSeq]

Other Designations BAF complex 53 kDa subunit|BAF53|BRG1-associated factor|INO80 complex subunit K|actin-related protein 4|hArpN beta