

ACVR1 rabbit monoclonal antibody

Catalog # H00000090-K Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human ACVR1 peptide using ARM Technology.
Immunogen	A synthetic peptide of human ACVR1 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human ACVR1 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — ACVR1

Entrez GeneID	90
GeneBank Accession#	ACVR1
Gene Name	ACVR1
Gene Alias	ACTRI, ACVR1A, ACVRLK2, ALK2, FOP, SKR1, TSRI
Gene Description	activin A receptor, type I
Omim ID	102576 135100
Gene Ontology	Hyperlink
Gene Summary	Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. This gene encodes activin A type I receptor which signals a particular transcriptional response in concert with activin type II receptors. Mutations in this gene are associated with fibrodysplasia ossificans progressive. [provided by RefSeq]
Other Designations	TGF-B superfamily receptor type II activin A receptor, type II-like kinase 2 activin A type I receptor hydroxyalkyl-protein kinase serine/threonine-protein kinase receptor R1

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [TGF-beta signaling pathway](#)

Disease

- [Abortion](#)
- [Breast Neoplasms](#)

- [Genetic Predisposition to Disease](#)
- [Infertility](#)
- [Intracranial Aneurysm](#)
- [Narcolepsy](#)
- [Neoplasms](#)
- [Obesity](#)
- [Ovarian cancer](#)
- [Ovarian Failure](#)
- [Ovarian Neoplasms](#)
- [Polycystic Ovary Syndrome](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Puberty](#)
- [Stroke](#)
- [Thrombophilia](#)
- [Tobacco Use Disorder](#)