


ACVR1B (Human) Matched Antibody Pair

Catalog # H00000091-AP42 Size 1 Set

Applications



Sandwich ELISA detection sensitivity ranging from 3 ng/ml to 100 ng/ml.

Specification

Product Description	This antibody pair set comes with a matched antibody pair to detect and quantify the protein level of human ACVR1B.
Reactivity	Human
Quality Control Testing	Standard curve using recombinant protein (H00000091-Q01) as an analyte. Sandwich ELISA detection sensitivity ranging from 3 ng/ml to 100 ng/ml.
Supplied Product	Antibody pair set content: 1. Capture antibody: mouse monoclonal anti-ACVR1B, IgG2a Kappa (100 ug) 2. Detection antibody: biotinylated mouse monoclonal anti-ACVR1B, IgG1 Kappa (50 ug) *Reagents are sufficient for at least 3-5 x 96 well plates using recommended protocols.
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

- ELISA Pair (Recombinant protein)

[Protocol Download](#)

Gene Info — ACVR1B

Entrez GeneID [91](#)

Gene Name ACVR1B

Gene Alias ACTRIB, ACVRLK4, ALK4, SKR2

Gene Description activin A receptor, type IB

Omim ID [601300](#)

Gene Ontology [Hyperlink](#)

Gene Summary

Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with a cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling, and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. This gene encodes activin A type IB receptor, composed of 11 exons. Alternative splicing and alternative polyadenylation result in 3 fully described transcript variants. The mRNA expression of variants 1, 2, and 3 is confirmed, and a potential fourth variant contains an alternative exon 8 and lacks exons 9 through 11, but its mRNA expression has not been confirmed. [provided by RefSeq]

Other Designations

activin A receptor, type II-like kinase 4|activin A type IB receptor|activin receptor-like kinase 4|serine(threonine) protein kinase receptor R2

Pathway

- [Adherens junction](#)
- [Chronic myeloid leukemia](#)
- [Colorectal cancer](#)
- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [MAPK signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)

- [TGF-beta signaling pathway](#)

Disease

- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Obesity](#)
- [Ovarian Failure](#)
- [Polycystic Ovary Syndrome](#)
- [Puberty](#)
- [Schizophrenia](#)
- [Thrombophilia](#)
- [Tobacco Use Disorder](#)