


ACVR1B monoclonal antibody (M01), clone 2D4

Catalog # H00000091-M01


Size 100 ug

Applications



Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged ACVR1B is approximately 0.03ng/ml as a capture antibody.



Western Blot detection against Immunogen (36.96 KDa) .

Specification

Product Description	Mouse monoclonal antibody raised against a partial recombinant ACVR1B.
Immunogen	ACVR1B (AAH00254, 24 a.a. ~ 126 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	SGPRGVQALLCACTSC LQANYTCETDGACMVSIFNLDGMEHHVRTCIPKVELVPAGKPFYCLSS EDLRNTHCCYTDYCNRIDLRVPSGHLKEPEHPSMWGPVE
Host	Mouse
Reactivity	Human

Interspecies Antigen Sequence

Mouse (93); Rat (95)

Isotype

IgG2a Kappa

Quality Control Testing

Antibody Reactive Against Recombinant Protein.
Western Blot detection against Immunogen (36.96 KDa) .

Storage Buffer

In 1x PBS, pH 7.4

Storage Instruction

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Recombinant protein)

[Protocol Download](#)

- Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged ACVR1B is approximately 0.03ng/ml as a capture antibody.

[Protocol Download](#)

- ELISA

Gene Info — ACVR1B

Entrez GeneID[91](#)**GeneBank Accession#**[BC000254](#)**Protein Accession#**[AAH00254](#)**Gene Name**

ACVR1B

Gene Alias

ACTRIB, ACVRLK4, ALK4, SKR2

Gene Description

activin A receptor, type IB

Omim ID[601300](#)**Gene Ontology**[Hyperlink](#)

Gene Summary

Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with a cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling, and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. This gene encodes activin A type IB receptor, composed of 11 exons. Alternative splicing and alternative polyadenylation result in 3 fully described transcript variants. The mRNA expression of variants 1, 2, and 3 is confirmed, and a potential fourth variant contains an alternative exon 8 and lacks exons 9 through 11, but its mRNA expression has not been confirmed. [provided by RefSeq]

Other Designations

activin A receptor, type II-like kinase 4|activin A type IB receptor|activin receptor-like kinase 4|serine(threonine) protein kinase receptor R2

Pathway

- [Adherens junction](#)
- [Chronic myeloid leukemia](#)
- [Colorectal cancer](#)
- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [MAPK signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [TGF-beta signaling pathway](#)

Disease

- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)

- [Obesity](#)
- [Ovarian Failure](#)
- [Polycystic Ovary Syndrome](#)
- [Puberty](#)
- [Schizophrenia](#)
- [Thrombophilia](#)
- [Tobacco Use Disorder](#)