


ALDH3A1 (Human) IP-WB Antibody Pair

Catalog # H00000218-PW1

Size 1 Set

Applications



Immunoprecipitation of ALDH3A1 transfected lysate using rabbit polyclonal anti-ALDH3A1 and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with mouse purified polyclonal anti-ALDH3A1.

Specification

Product Description	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.
Reactivity	Human
Quality Control Testing	Immunoprecipitation-Western Blot (IP-WB) Immunoprecipitation of ALDH3A1 transfected lysate using rabbit polyclonal anti-ALDH3A1 and Protein A Magnetic Bead (U0007), and immunoblotted with mouse purified polyclonal anti-ALDH3A1.
Supplied Product	Antibody pair set content: 1. Antibody pair for IP: rabbit polyclonal anti-ALDH3A1 (300 ul) 2. Antibody pair for WB: mouse purified polyclonal anti-ALDH3A1 (50 ug)
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

- Immunoprecipitation-Western Blot

[Protocol Download](#)

Gene Info — ALDH3A1

Entrez GeneID [218](#)

Gene Name ALDH3A1

Gene Alias ALDH3, ALDHIII, MGC10406

Gene Description aldehyde dehydrogenase 3 family, memberA1

Omim ID [100660](#)

Gene Ontology [Hyperlink](#)

Gene Summary Aldehyde dehydrogenases oxidize various aldehydes to the corresponding acids. They are involved in the detoxification of alcohol-derived acetaldehyde and in the metabolism of corticosteroids, biogenic amines, neurotransmitters, and lipid peroxidation. The enzyme encoded by this gene forms a cytoplasmic homodimer that preferentially oxidizes aromatic and medium-chain (6 carbons or more) saturated and unsaturated aldehyde substrates. It is thought to promote resistance to UV and 4-hydroxy-2-nonenal-induced oxidative damage in the cornea. The gene is located within the Smith-Magenis syndrome region on chromosome 17. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq]

Other Designations aldehyde dehydrogenase 3A1|aldehyde dehydrogenase isozyme 3|aldehyde dehydrogenase type III|aldehyde dehydrogenase, dimeric NADP-preferring|stomach aldehyde dehydrogenase

Pathway

- [Drug metabolism - cytochrome P450](#)
- [Glycolysis / Gluconeogenesis](#)
- [Histidine metabolism](#)
- [Metabolic pathways](#)
- [Metabolism of xenobiotics by cytochrome P450](#)
- [Phenylalanine metabolism](#)
- [Tyrosine metabolism](#)

Disease

- [Attention Deficit Disorder with Hyperactivity](#)

- [Autistic Disorder](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Drug Toxicity](#)
- [Genetic Predisposition to Disease](#)
- [Kidney Failure](#)
- [NARP](#)
- [Neoplasms](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Schizophrenia](#)
- [Tobacco Use Disorder](#)