

BIRC3 rabbit monoclonal antibody

Catalog # H00000330-K Size 100 ug x up to 3

| Specification | |
|-------------------------|---|
| Product Description | Rabbit monoclonal antibody raised against a human BIRC3 peptide using ARM Technology. |
| Immunogen | A synthetic peptide of human BIRC3 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence. |
| Host | Rabbit |
| Library Construction | Non-fusion antibody library from rabbit spleen (<u>ARM Technology</u>). |
| Expression | Overexpression vector and transfection into 293H cell line. |
| Reactivity | Human |
| Purification | Protein A |
| Isotype | lgG |
| Quality Control Testing | Antibody reactive against human BIRC3 peptide by ELISA and mammalian transfected lysate by We stern Blot. |
| Storage Buffer | In 1x PBS, pH 7.4 |
| Storage Instruction | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing. |
| Deliverable | Up to three rabbit lgG clones of 100 ug each will be delivered to customer. |
| Note | Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request. |

Applications

Western Blot (Transfected lysate)

Protocol Download



ELISA

| Gene Info — BIRC3 | |
|---------------------|--|
| Entrez GenelD | 330 |
| GeneBank Accession# | BIRC3 |
| Gene Name | BIRC3 |
| Gene Alias | AIP1, API2, CIAP2, HAIP1, HIAP1, MALT2, MIHC, RNF49 |
| Gene Description | baculoviral IAP repeat-containing 3 |
| Omim ID | 601721 |
| Gene Ontology | <u>Hyperlink</u> |
| Gene Summary | The protein encoded by this gene is a member of a family of proteins that inhibits apoptosis by bi nding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interf ering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by ser um deprivation but does not affect apoptosis resulting from exposure to menadione, a potent indu cer of free radicals. The amino acid sequence predicts three baculovirus IAP repeat domains and a ring finger domain. Transcript variants encoding the same isoform have been identified. [provid ed by RefSeq |
| Other Designations | TNFR2-TRAF signaling complex protein apoptosis inhibitor 2 baculoviral IAP repeat-containing protein 3 inhibitor of apoptosis protein 1 mammalian IAP homolog C |

Pathway

- Apoptosis
- Focal adhesion
- Pathways in cancer
- Small cell lung cancer
- <u>Ubiquitin mediated proteolysis</u>

Disease

Adenocarcinoma



- Cardiovascular Diseases
- Diabetes Mellitus
- Ductus Arteriosus
- Edema
- Esophageal Neoplasms
- Genetic Predisposition to Disease
- Infant
- Lung Neoplasms
- Pulmonary Disease
- Urinary Bladder Neoplasms
- Werner syndrome