


MaxPab®

APRT purified MaxPab rabbit polyclonal antibody (D01P)

Catalog # H00000353-D01P


Size 100 ug

Applications



Western Blot (Cell lysate)

APRT MaxPab rabbit polyclonal antibody. Western Blot analysis of APRT expression in K-562.



Western Blot (Transfected lysate)

Western Blot analysis of APRT expression in transfected 293T cell line ([H00000353-T01](#)) by APRT MaxPab polyclonal antibody.

Lane 1: APRT transfected lysate(19.60 KDa).

Lane 2: Non-transfected lysate.

Specification

Product Description	Rabbit polyclonal antibody raised against a full-length human APRT protein.
Immunogen	APRT (NP_000476.1, 1 a.a. ~ 180 a.a) full-length human protein.
Sequence	MADSELQLVEQRIRSFDPFPTPGVVFRDISPVLKDPASFRAAIGLLARHLKATHGGRIDYIAGLDSR GFLFGPSLAQELGLGCVLIRKRGKLPGPTLWASYSLEYGKAELEIQKDALEPGQRVVVVDDLLAT GGTMNAACELLGRLQAEVLECVSLVELTSLKGREKLAPVPFFSLLQYE
Host	Rabbit
Reactivity	Human

Interspecies Antigen Sequence	Mouse (83); Rat (87)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Cell lysate)

APRT MaxPab rabbit polyclonal antibody. Western Blot analysis of APRT expression in K-562.

[Protocol Download](#)

- Western Blot (Transfected lysate)

Western Blot analysis of APRT expression in transfected 293T cell line ([H00000353-T01](#)) by APRT MaxPab polyclonal antibody.

Lane 1: APRT transfected lysate(19.60 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

Gene Info — APRT

Entrez GeneID	353
GeneBank Accession#	NM_000485.2
Protein Accession#	NP_000476.1
Gene Name	APRT
Gene Alias	AMP, DKFZp686D13177, MGC125856, MGC125857, MGC129961
Gene Description	adenine phosphoribosyltransferase
Omim ID	102600
Gene Ontology	Hyperlink

Gene Summary

Adenine phosphoribosyltransferase belongs to the purine/pyrimidine phosphoribosyltransferase family. A conserved feature of this gene is the distribution of CpG dinucleotides. This enzyme catalyzes the formation of AMP and inorganic pyrophosphate from adenine and 5-phosphoribosyl-1-pyrophosphate (PRPP). It also produces adenine as a by-product of the polyamine biosynthesis pathway. A homozygous deficiency in this enzyme causes 2,8-dihydroxyadenine urolithiasis. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations

AMP diphosphorylase|AMP pyrophosphorylase|adenine phosphoribosyltransferase, isoform a|transphosphoribosidase

Pathway

- [Metabolic pathways](#)
- [Purine metabolism](#)