


ASGR2 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # H00000433-T03


Size 100 uL

Applications



SDS-PAGE Gel

ASGR2 transfected lysate.



Western Blot

Lane 1: ASGR2 transfected lysate (32.6 KDa)

Lane 2: Non-transfected lysate.

Specification

Transfected Cell Line 293T

Plasmid pCMV-ASGR2 full-length

Host Human

Theoretical MW (kDa) 32.6

Quality Control Testing Transient overexpression cell lysate was tested with Anti-ASGR2 antibody ([H00000433-B02P](#)) by Western Blots.
SDS-PAGE Gel
ASGR2 transfected lysate.
Western Blot
Lane 1: ASGR2 transfected lysate (32.6 KDa)
Lane 2: Non-transfected lysate.

Storage Buffer

1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot

Gene Info — ASGR2

Entrez GeneID[433](#)**GeneBank Accession#**[NM_080913](#)**Protein Accession#**[NP_550435.1](#)**Gene Name**

ASGR2

Gene Alias

ASGP-R, CLEC4H2, Hs.1259, L-H2

Gene Description

asialoglycoprotein receptor 2

Omim ID[108361](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This cell surface receptor binds to galactose-terminated glycoproteins. It transports these glycoproteins via a series of membrane vesicles and tubules to an acidic-sorting organelle where the receptor and ligand dissociates. Then the receptor is recycled back to the cell surface. There are four alternatively spliced transcript variants of this gene. [provided by RefSeq]

Other Designations

C-type lectin domain family 4, member H2|OTTHUMP00000128352|OTTHUMP00000128353|asialoglycoprotein receptor H2|hepatic lectin H2