

# ASPH rabbit monoclonal antibody

Catalog # H00000444-K

Size 100 ug x up to 3

## Specification

<b>Product Description</b>	Rabbit monoclonal antibody raised against a human ASPH peptide using ARM Technology.
<b>Immunogen</b>	A synthetic peptide of human ASPH is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
<b>Host</b>	Rabbit
<b>Library Construction</b>	Non-fusion antibody library from rabbit spleen ( <a href="#">ARM Technology</a> ).
<b>Expression</b>	Overexpression vector and transfection into 293H cell line.
<b>Reactivity</b>	Human
<b>Purification</b>	Protein A
<b>Isotype</b>	IgG
<b>Quality Control Testing</b>	Antibody reactive against human ASPH peptide by ELISA and mammalian transfected lysate by Western Blot.
<b>Storage Buffer</b>	In 1x PBS, pH 7.4
<b>Storage Instruction</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Deliverable</b>	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
<b>Note</b>	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) <sub>2</sub> , IgG, scFv and different Fc and non-Fc conjugates per customer request.

## Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

## Gene Info — ASPH

Entrez GeneID [444](#)

GeneBank Accession# [ASPH](#)

Gene Name ASPH

Gene Alias BAH, CASQ2BP1, HAAH, JCTN, junctin

Gene Description aspartate beta-hydroxylase

Omim ID [600582](#)

Gene Ontology [Hyperlink](#)

**Gene Summary**

This gene is thought to play an important role in calcium homeostasis. The gene is expressed from two promoters and undergoes extensive alternative splicing. The encoded set of proteins share varying amounts of overlap near their N-termini but have substantial variations in their C-terminal domains resulting in distinct functional properties. The longest isoforms (a and f) include a C-terminal Aspartyl/Asparaginyl beta-hydroxylase domain that hydroxylates aspartic acid or asparagine residues in the epidermal growth factor (EGF)-like domains of some proteins, including protein C, coagulation factors VII, IX, and X, and the complement factors C1R and C1S. Other isoforms differ primarily in the C-terminal sequence and lack the hydroxylase domain, and some have been localized to the endoplasmic and sarcoplasmic reticulum. Some of these isoforms are found in complexes with calsequestrin, triadin, and the ryanodine receptor, and have been shown to regulate calcium release from the sarcoplasmic reticulum. Some isoforms have been implicated in metastasis. [provided by RefSeq]

**Other Designations** aspartyl/asparaginyl-beta-hydroxylase|humbug|junctate|peptide-aspartate beta-dioxygenase