


Full-Length

ATF4 (Human) Recombinant Protein (P02)

Catalog # H00000468-P02

Size 25 ug, 10 ug

Applications



Specification

Product Description

Human ATF4 full-length ORF (AAH16855, 1 a.a. - 351 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence

MTEMSFLSSEVLVGDLMSPFDQSGLGAEESLGLDDYLEVAKHFKPHGFSSDKAKAGSSEWLA
VDGLVSPSNNKEDAFSGTDWMLEKMDLKEFDLDALLGIDDLTMPDDLTTLDDTCDLFAPLV
QETNKQPPQTVNPIGHLPESLTKPDQVAPFTFLQPLPLSPGVLSSTPDHSFSLELGSEVDITEGDR
KPDYTA YVAMIPQCIKEEDTPSDNDSGICMSPESYLGSPQHSPSTRGSPNRSPLSPGVLGCSARP
KPYDPPGEKMVA AKVKGEKLDKCLKKMEQNKTAATRYRQKKRAEQEALTGECKELEKKNEALK
ERADSLAKEIQYLKDLIEEVRKARGKKRVP

Host

Wheat Germ (in vitro)

Theoretical MW (kDa)

64.35

Preparation Method

[in vitro wheat germ expression system](#)

Purification

Glutathione Sepharose 4 Fast Flow

Quality Control Testing

12.5% SDS-PAGE Stained with Coomassie Blue.

Storage Buffer

50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Note

Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — ATF4

Entrez GeneID [468](#)

GeneBank Accession# [BC016855](#)

Protein Accession# [AAH16855](#)

Gene Name ATF4

Gene Alias CREB-2, CREB2, TAXREB67, TXREB

Gene Description activating transcription factor 4 (tax-responsive enhancer element B67)

Omim ID [604064](#)

Gene Ontology [Hyperlink](#)

Gene Summary

This gene encodes a transcription factor that was originally identified as a widely expressed mammalian DNA binding protein that could bind a tax-responsive enhancer element in the LTR of HTLV-1. The encoded protein was also isolated and characterized as the cAMP-response element binding protein 2 (CREB-2). The protein encoded by this gene belongs to a family of DNA-binding proteins that includes the AP-1 family of transcription factors, cAMP-response element binding proteins (CREBs) and CREB-like proteins. These transcription factors share a leucine zipper region that is involved in protein-protein interactions, located C-terminal to a stretch of basic amino acids that functions as a DNA binding domain. Two alternative transcripts encoding the same protein have been described. Two pseudogenes are located on the X chromosome at q28 in a region containing a large inverted duplication. [provided by RefSeq]

Other Designations activating transcription factor 4|cAMP response element-binding protein 2

Pathway

- [GnRH signaling pathway](#)
- [Long-term potentiation](#)
- [MAPK signaling pathway](#)
- [Neurotrophin signaling pathway](#)
- [Prostate cancer](#)

Disease

- [Bipolar Disorder](#)
- [Genetic Predisposition to Disease](#)
- [Mental Disorders](#)
- [Neuropsychological Tests](#)
- [Schizophrenia](#)