


# ATP6V1G2 monoclonal antibody (M02), clone 2E11

Catalog # H00000534-M02


Size 100 ug

## Applications




### Western Blot (Cell lysate)

ATP6V1G2 monoclonal antibody (M02), clone 2E11 Western Blot analysis of ATP6V1G2 expression in HepG2 ( Cat # L019V1 ).




### Western Blot (Cell lysate)

ATP6V1G2 monoclonal antibody (M02), clone 2E11. Western Blot analysis of ATP6V1G2 expression in Raw 264.7 ( Cat # L024V1 ).




### Western Blot (Cell lysate)

ATP6V1G2 monoclonal antibody (M02), clone 2E11. Western Blot analysis of ATP6V1G2 expression in PC-12 ( Cat # L012V1 ).



### Western Blot (Cell lysate)

ATP6V1G2 monoclonal antibody (M02), clone 2E11. Western Blot analysis of ATP6V1G2 expression in NIH/3T3 ( Cat # L018V1 ).




## Western Blot (Transfected lysate)

Western Blot analysis of ATP6V1G2 expression in transfected 293T cell line by ATP6V1G2 monoclonal antibody (M02), clone 2E11.

Lane 1: ATP6V1G2 transfected lysate(13.6 KDa).

Lane 2: Non-transfected lysate.



Western Blot detection against Immunogen (34.32 KDa).

## Specification

<b>Product Description</b>	Mouse monoclonal antibody raised against a partial recombinant ATP6V1G2.
<b>Immunogen</b>	ATP6V1G2 (NP_569730, 41 a.a. ~ 118 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
<b>Sequence</b>	QMEVEQYRREREHEFQSKQQAAMGSQGNLSAEVEQATRRQVQGMQSSQQRNRERVLAQLLG MVCDVRPQVHPNYRISA
<b>Host</b>	Mouse
<b>Reactivity</b>	Human, Mouse, Rat
<b>Isotype</b>	IgG2b Lambda
<b>Quality Control Testing</b>	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (34.32 KDa).
<b>Storage Buffer</b>	In 1x PBS, pH 7.4
<b>Storage Instruction</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

## Applications

- Western Blot (Cell lysate)

ATP6V1G2 monoclonal antibody (M02), clone 2E11 Western Blot analysis of ATP6V1G2 expression in HepG2 ( Cat # L019V1 ).

[Protocol Download](#)

- Western Blot (Cell lysate)

ATP6V1G2 monoclonal antibody (M02), clone 2E11. Western Blot analysis of ATP6V1G2 expression in Raw 264.7 ( Cat # L024V1 ).

[Protocol Download](#)

- Western Blot (Cell lysate)

ATP6V1G2 monoclonal antibody (M02), clone 2E11. Western Blot analysis of ATP6V1G2 expression in PC-12 ( Cat # L012V1 ).

[Protocol Download](#)

- Western Blot (Cell lysate)

ATP6V1G2 monoclonal antibody (M02), clone 2E11. Western Blot analysis of ATP6V1G2 expression in NIH/3T3 ( Cat # L018V1 ).

[Protocol Download](#)

- Western Blot (Transfected lysate)

Western Blot analysis of ATP6V1G2 expression in transfected 293T cell line by ATP6V1G2 monoclonal antibody (M02), clone 2E11.

Lane 1: ATP6V1G2 transfected lysate(13.6 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

- Western Blot (Recombinant protein)

[Protocol Download](#)

- ELISA

## Gene Info — ATP6V1G2

Entrez GeneID

[534](#)

GeneBank Accession#

[NM\\_130463](#)

Protein Accession#	<a href="#">NP_569730</a>
Gene Name	ATP6V1G2
Gene Alias	ATP6G, ATP6G2, NG38, VMA10
Gene Description	ATPase, H <sup>+</sup> transporting, lysosomal 13kDa, V1 subunit G2
Omim ID	<a href="#">606853</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	<p>This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of intracellular compartments of eukaryotic cells. V-ATPase dependent acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is one of three V1 domain G subunit proteins. This gene had previous gene symbols of ATP6G and ATP6G2. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq]</p>
Other Designations	ATPase, H <sup>+</sup> transporting, lysosomal (vacuolar proton pump) subunit G ATPase, H <sup>+</sup> transporting, lysosomal, V1 subunit G2 H <sup>+</sup> -transporting two-sector ATPase, subunit G2 OTTHUMP00000029286 OTTHUMP00000036058 OTTHUMP00000036060 V-ATPase 13 kDa subunit 2 V-ATPa

## Pathway

- [Epithelial cell signaling in Helicobacter pylori infection](#)
- [Metabolic pathways](#)
- [Oxidative phosphorylation](#)
- [Vibrio cholerae infection](#)

## Disease

- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)

- [Lupus Erythematosus](#)
- [Malaria](#)
- [Tobacco Use Disorder](#)