


Full-Length

BAD (Human) Recombinant Protein (P01)

Catalog # H00000572-P01

Size 25 ug, 10 ug

Applications



Specification

Product Description

Human BAD full-length ORF (AAH01901.1, 1 a.a. - 168 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence

MFQIPEFEPSEQEDSSSAERGLGPSPAGDGPSGSGKHHRQAPGLLWDASHQQEQPTSSSHHG
GAGAVEIRSRHSSYPAGTEDDEGMGEESPFRGRSRSAPPNLWAAQRYGRELRRMSDEFVDSF
KKGLPRPKSAGTATQMRQSSSWTRVFQSWWDRNLGRGSSAPSQ

Host

Wheat Germ (in vitro)

Theoretical MW (kDa)

44.11

Preparation Method

[in vitro wheat germ expression system](#)

Purification

Glutathione Sepharose 4 Fast Flow

Quality Control Testing

12.5% SDS-PAGE Stained with Coomassie Blue.

Storage Buffer

50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Note

Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — BAD

Entrez GeneID [572](#)

GeneBank Accession# [BC001901.1](#)

Protein Accession# [AAH01901.1](#)

Gene Name BAD

Gene Alias BBC2, BCL2L8

Gene Description BCL2-associated agonist of cell death

Omim ID [603167](#)

Gene Ontology [Hyperlink](#)

Gene Summary The protein encoded by this gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. [provided by RefSeq]

Other Designations BCL-X/BCL-2 binding protein|BCL2-antagonist of cell death protein|BCL2-binding component 6|BCL2-binding protein

Pathway

- [Acute myeloid leukemia](#)

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Chronic myeloid leukemia](#)
- [Colorectal cancer](#)
- [Endometrial cancer](#)
- [ErbB signaling pathway](#)
- [Focal adhesion](#)
- [Insulin signaling pathway](#)
- [Melanoma](#)
- [Neurotrophin signaling pathway](#)
- [Non-small cell lung cancer](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [VEGF signaling pathway](#)

Disease

- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Lymphoma](#)
- [Parkinson disease](#)
- [Thyroid Neoplasms](#)