


MaxPab®

BCL3 purified MaxPab mouse polyclonal antibody (B01P)

Catalog # H00000602-B01P

Size 50 ug

Applications




Western Blot (Transfected lysate)

Western Blot analysis of BCL3 expression in transfected 293T cell line ([H00000602-T01](#)) by BCL3 MaxPab polyclonal antibody.

Lane 1: BCL3 transfected lysate(49.06 KDa).

Lane 2: Non-transfected lysate.



Immunofluorescence

Immunofluorescence of purified MaxPab antibody to BCL3 on HeLa cell. [antibody concentration 10 ug/ml]

Specification

Product Description

Mouse polyclonal antibody raised against a full-length human BCL3 protein.

Immunogen

BCL3 (NP_005169.1, 1 a.a. ~ 446 a.a) full-length human protein.

Sequence

MDEGPVDLRTRPKAAGLPGAALPLRKRLRAPSPEPAAPRGAAGLVVPLDPLRGGCDLPAVPG
PPHGLARPEALYYPGALLPLYPTRAMGSPFPLVNLPTLYPMMCPMEHPLSADIAMATRADEDG
DTPLHIAVVQGNLPAVHRLVNLFFQQGGRELDIYNNLRQTPLHLAVITTLPSVVRLLVTAGASPMALD
RHGQTAAHLACEHRSPTCLRALLDSAAPGTLDLEARNYDGLTALHVAVNTECQETVQLLLERGA
DIDAVDIKSGRSLIHAVENNSLSMVQLLLQHGANVNAQMYSGSSALHSASGRGLPLVRTLVRS
GADSSLKNCHNDTPLMVARSRVIDILRGKATRPASTSQPDPSPPDRSANTSPESSRLSSNGLLS
ASPSSSPSQSPPRDPPGFPMAPPNFFLPSPSPPAFLPFAGVLRGPRPVPPSPAPGGS

Host	Mouse
Reactivity	Human
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of BCL3 expression in transfected 293T cell line ([H00000602-T01](#)) by BCL3 MaxPab polyclonal antibody.

Lane 1: BCL3 transfected lysate(49.06 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

- Immunofluorescence

Immunofluorescence of purified MaxPab antibody to BCL3 on HeLa cell. [antibody concentration 10 ug/ml]

Gene Info — BCL3

Entrez GeneID	602
GeneBank Accession#	NM_005178
Protein Accession#	NP_005169.1
Gene Name	BCL3
Gene Alias	BCL4, D19S37
Gene Description	B-cell CLL/lymphoma 3
Omim ID	109560
Gene Ontology	Hyperlink

Gene Summary

This gene is a proto-oncogene candidate. It is identified by its translocation into the immunoglobulin alpha-locus in some cases of B-cell leukemia. The protein encoded by this gene contains seven ankyrin repeats, which are most closely related to those found in I kappa B proteins. This protein functions as a transcriptional co-activator that activates through its association with NF-kappa B homodimers. The expression of this gene can be induced by NF-kappa B, which forms a part of the autoregulatory loop that controls the nuclear residence of p50 NF-kappa B. [provided by RefSeq]

Other Designations

B-cell leukemia/lymphoma 3|B-cell lymphoma 3-encoded protein|chronic lymphatic leukemia protein

Disease

- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Crohn Disease](#)
- [Disease Models](#)
- [Genetic Predisposition to Disease](#)