

BMP5 rabbit monoclonal antibody

Catalog # H00000653-K Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human BMP5 peptide using ARM Technology.
Immunogen	A synthetic peptide of human BMP5 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (<u>ARM Technology</u>).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	lgG
Quality Control Testing	Antibody reactive against human BMP5 peptide by ELISA and mammalian transfected lysate by We stern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit lgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

Western Blot (Transfected lysate)

Protocol Download



ELISA

Gene Info — BMP5	
Entrez GenelD	<u>653</u>
GeneBank Accession#	BMP5
Gene Name	BMP5
Gene Alias	MGC34244
Gene Description	bone morphogenetic protein 5
Omim ID	112265
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene encodes a member of the bone morphogenetic protein family which is part of the transf orming growth factor-beta superfamily. The superfamily includes large families of growth and diffe rentiation factors. Bone morphogenetic proteins were originally identified by an ability of deminer alized bone extract to induce endochondral osteogenesis in vivo in an extraskeletal site. These proteins are synthesized as prepropeptides, cleaved, and then processed into dimeric proteins. This protein may act as an important signaling molecule within the trabecular meshwork and optic nerve head, and may play a potential role in glaucoma pathogenesis. This gene is differentially regulated during the formation of various tumors. [provided by RefSeq
Other Designations	OTTHUMP0000016650

Pathway

- Cytokine-cytokine receptor interaction
- Hedgehog signaling pathway
- TGF-beta signaling pathway

Disease

- Congenital Abnormalities
- Genetic Predisposition to Disease
- Obesity



- Osteoarthritis
- Ovarian Failure
- Polycystic Ovary Syndrome
- Puberty
- Thrombophilia
- Tobacco Use Disorder