


MaxPab®

BYSL purified MaxPab mouse polyclonal antibody (B01P)


Catalog # H00000705-B01P Size 50 ug

Applications



Western Blot (Cell lysate)

BYSL MaxPab polyclonal antibody. Western Blot analysis of BYSL expression in Jurkat




Western Blot (Transfected lysate)

Western Blot analysis of BYSL expression in transfected 293T cell line (<u>H00000705-T01</u>) by BYSL MaxPab polyclonal antibody.

Lane 1: BYSL transfected lysate(35.75 KDa).

Lane 2: Non-transfected lysate.



Immunofluorescence

Immunofluorescence of <u>purified</u> MaxPab antibody to BYSL on HeLa cell. [antibody concentration 10 ug/ml]

Specification

Product Description

Mouse polyclonal antibody raised against a full-length human BYSL protein.



Product Information

Immunogen	BYSL (AAH07340, 1 a.a. ~ 324 a.a) full-length human protein.
Sequence	MTAAGHHAEVVVDPEDERAIEMFMNKNPPARRTLADIIMEKLTEKQTEVETVMSEVSGFPMPQL DPRVLEVYRGVREVLSKYRSGKLPKAFKIIPALSNWDQILYVTEPEAWTAAAMYQATRIFASNLKE RMAQRFYNLVLLPRVRDDVAEYKRLNFHLYMALKKALFKPGAWFKGILIPLCESGTCTLREAIIVGSI ITKCSIPVLHSSAAMLKIAEMEYSGANSIFLRLLLDKKYALPYRVLDALVFHFLGFRTEKRELPVLWH QCLLTLVQRYKADLATDQKEALLELLRLQPHPQLSPEIRRELQSAVPRDVEDVPITVE
Host	Mouse
Reactivity	Human
Interspecies Antigen Sequence	Mouse (94); Rat (95)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

Western Blot (Cell lysate)

BYSL MaxPab polyclonal antibody. Western Blot analysis of BYSL expression in Jurkat.

Protocol Download

Western Blot (Transfected lysate)

 $We stern \ Blot \ analysis \ of \ BYSL \ expression \ in \ transfected \ 293T \ cell \ line \ (\underline{H00000705\text{-}T01}) \ by \ BYSL \ MaxPab \ polyclonal \ antibody.$

Lane 1: BYSL transfected lysate(35.75 KDa).

Lane 2: Non-transfected lysate.

Protocol Download

Immunofluorescence

Immunofluorescence of <u>purified</u> MaxPab antibody to BYSL on HeLa cell. [antibody concentration 10 ug/ml]

Gene		DVCI
(-AnA	INTO —	\bowtie \vee \vee

Entrez GeneID	<u>705</u>
GeneBank Accession#	BC007340



Product Information

Protein Accession#	<u>AAH07340</u>
Gene Name	BYSL
Gene Alias	BYSTIN
Gene Description	bystin-like
Omim ID	603871
Gene Ontology	<u>Hyperlink</u>
Gene Summary	Bystin is expressed as a 2-kb major transcript and a 3.6-kb minor transcript in SNG-M cells and in human trophoblastic teratocarcinoma HT-H cells. Protein binding assays determined that bystin binds directly to trophinin and tastin, and that binding is enhanced when cytokeratins 8 and 18 are present. Immunocytochemistry of HT-H cells showed that bystin colocalizes with trophinin, tastin, and the cytokeratins, suggesting that these molecules form a complex in trophectoderm cells at the time of implantation. Using immunohistochemistry it was determined that trophinin and bystin are found in the placenta from the sixth week of pregnancy. Both proteins were localized in the cytoplas mof the syncytiotrophoblast in the chorionic villi and in endometrial decidual cells at the uteroplac ental interface. After week 10, the levels of trophinin, tastin, and bystin decreased and then disappeared from placental villi. [provided by RefSeq
Other Designations	OTTHUMP00000039776 by the ribosomal protein s6 gene, drosophila, homolog-like bystin

Disease

• Kidney Failure