


CCNC 293T Cell Transient Overexpression Lysate(Denatured)


Catalog # H00000892-T04 Size 100 uL

Applications



SDS-PAGE Gel

CCNC transfected lysate.



Western Blot

Lane 1: CCNC transfected lysate (31.24 KDa)

Lane 2: Non-transfected lysate.

Specification	
Transfected Cell Line	293T
Plasmid	pCMV-CCNC full-length
Host	Human
Theoretical MW (kDa)	31.24
Interspecies Antigen Sequence	Mouse (99); Rat (99)



Product Information

estern Blots. SDS-PAGE Gel CCNC transfected lysate. Western Blot Lane 1: CCNC transfected lysate (31.24 KDa)	
Lane 2: Non-transfected lysate.	
1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bro mophenol blue)	
Store at -80°C. Aliquot to avoid repeated freezing and thawing.	

Applications

Western Blot

Gene Info — CCNC	
Entrez GeneID	<u>892</u>
GeneBank Accession#	BC056153.1
Protein Accession#	AAH56153.1
Gene Name	CCNC
Gene Alias	CycC
Gene Description	cyclin C
Omim ID	<u>123838</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a member of the cyclin family of proteins. The encoded protein n interacts with cyclin-dependent kinase 8 and induces the phophorylation of the carboxy-terminal domain of the large subunit of RNA polymerase II. The level of mRNAs for this gene peaks in the G1 phase of the cell cycle. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq
Other Designations	OTTHUMP00000016897