

CD1B rabbit monoclonal antibody

Catalog # H00000910-K Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human CD1B peptide using ARM Technology.
Immunogen	A synthetic peptide of human CD1B is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human CD1B peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	<ol style="list-style-type: none"> Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab)₂, IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — CD1B

Entrez GeneID	910
GeneBank Accession#	CD1B
Gene Name	CD1B
Gene Alias	CD1, CD1A, MGC125990, MGC125991, R1
Gene Description	CD1b molecule
Omim ID	188360
Gene Ontology	Hyperlink
Gene Summary	<p>This gene encodes a member of the CD1 family of transmembrane glycoproteins, which are structurally related to the major histocompatibility complex (MHC) proteins and form heterodimers with beta-2-microglobulin. The CD1 proteins mediate the presentation of primarily lipid and glycolipid antigens of self or microbial origin to T cells. The human genome contains five CD1 family genes organized in a cluster on chromosome 1. The CD1 family members are thought to differ in their cellular localization and specificity for particular lipid ligands. The protein encoded by this gene localizes to late endosomes and lysosomes via a tyrosine-based motif in the cytoplasmic tail, and requires vesicular acidification to bind lipid antigens. [provided by RefSeq]</p>
Other Designations	CD1B antigen CD1B antigen, b polypeptide OTTHUMP00000018908 T-cell surface glycoprotein CD1b cortical thymocyte antigen CD1B differentiation antigen CD1-alpha-3

Pathway

- [Hematopoietic cell lineage](#)

Disease

- [Lung Diseases](#)
- [Mycobacterium Infections](#)