


MaxPab®

CD86 purified MaxPab mouse polyclonal antibody (B01P)

Catalog # H00000942-B01P

Size 50 ug

Applications



Western Blot (Transfected lysate)

Western Blot analysis of CD86 expression in transfected 293T cell line ([H00000942-T01](#)) by CD86 MaxPab polyclonal antibody.

Lane 1: CD86 transfected lysate(36.19 KDa).

Lane 2: Non-transfected lysate.

Specification

Product Description	Mouse polyclonal antibody raised against a full-length human CD86 protein.
Immunogen	CD86 (AAH40261.1, 1 a.a. ~ 329 a.a) full-length human protein.
Sequence	MDPQCTMGLSNILFVMAFLLSGAAPLKIQA YFNETADLPCQFANSQNQSLSELVFWQDQENLVL NEVYLGKEKFD SVHSKYMGR TSFDSWTLRLHNLQIKDKGLYQCIIHHKKPTGMIRIHQMNSELS VLANSQPEMPISNITENVYINLTCSSIHGYPEPKM SVLLRTKNSTIEYDGMQKSQDNVTELYDVSI SLSVSFPDVT SNMTIFCILETDKTRLLSSPFSIELEDPPQPPDHIPWITAVLPTVIICVMVFCLILWKW KKKKRPRNSYKCGTNTMEREESEQTKKREKIHIPERSDETQRVFKSSKTSSCDKSDTCF
Host	Mouse
Reactivity	Human
Interspecies Antigen Sequence	Mouse (50); Rat (57)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of CD86 expression in transfected 293T cell line ([H00000942-T01](#)) by CD86 MaxPab polyclonal antibody.

Lane 1: CD86 transfected lysate(36.19 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

Gene Info — CD86

Entrez GeneID [942](#)

GeneBank Accession# [BC040261.1](#)

Protein Accession# [AAH40261.1](#)

Gene Name CD86

Gene Alias B7-2, B70, CD28LG2, LAB72, MGC34413

Gene Description CD86 molecule

Omim ID [601020](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in two transcript variants encoding different isoforms. Additional transcript variants have been described, but their full-length sequences have not been determined. [provided by RefSeq]

Other Designations B-lymphocyte activation antigen B7-2|B7-2 antigen|CD28 antigen ligand 2|CD86 antigen|CD86 antigen (CD28 antigen ligand 2, B7-2 antigen)|CTLA-4 counter-receptor B7.2

Pathway

- [Allograft rejection](#)

- [Autoimmune thyroid disease](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Graft-versus-host disease](#)
- [Systemic lupus erythematosus](#)
- [Toll-like receptor signaling pathway](#)
- [Type I diabetes mellitus](#)

Disease

- [Acute Disease](#)
- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Bronchiolitis](#)
- [Cardiovascular Diseases](#)
- [Chronic Disease](#)
- [Colorectal Neoplasms](#)
- [Coronary Artery Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Drug Eruptions](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Hepatitis B](#)
- [Hepatitis C](#)
- [Hypersensitivity](#)
- [Infant](#)

- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Lupus Erythematosus](#)
- [Malignant melanoma](#)
- [Melanoma](#)
- [Multiple Sclerosis](#)
- [Papillomavirus Infections](#)
- [Pemphigus](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)
- [Sarcoidosis](#)
- [Scleroderma](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Werner syndrome](#)