

Proteoliposomes

Full-Length

TNFSF8 (Human) Recombinant Protein

Catalog # H00000944-G01

Size 10 ug

Specification

Product Description	Human TNFSF8 full-length ORF (NP_001235.1) recombinant protein without tag. This product is belong to Proteoliposome (PL).
Sequence	MDPGLQQALNGMAPPGDTAMHVPAGSVASHLGTTSRSYFYLTATLALCLVFTVATIMVLVVQRT DSIPNSPDNVPLKGGNCSEDLICILKRAPFKKSWAYLQVAKHLNKTLSWNKDGLHGVRVYQDGN LVIQFPGLYFIICQLQFLVQCPNNSVDLKLELLINKHIKKQALVTVCESGMQTKHVVYQNLSSQFLLDYL QVNTTISVNVDTFQYIDTSTFPLENVLSIFLYSNSD
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	26
Interspecies Antigen Sequence	Mouse (70)
Form	Liquid
Preparation Method	in vitro wheat germ expression system with proprietary liposome technology
Purification	None
Recommend Usage	Heating may cause protein aggregation. Please do not heat this product before electrophoresis.
Storage Buffer	25 mM Tris-HCl of pH8.0 containing 2% glycerol.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

Applications

- Antibody Production

Gene Info — TNFSF8

Entrez GeneID [944](#)

GeneBank Accession# [NM_001244.2](#)

Protein Accession# [NP_001235.1](#)

Gene Name TNFSF8

Gene Alias CD153, CD30L, CD30LG, MGC138144

Gene Description tumor necrosis factor (ligand) superfamily, member 8

Omim ID [603875](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF8/CD30, which is a cell surface antigen and a marker for Hodgkin lymphoma and related hematologic malignancies. The engagement of this cytokine expressed on B cell surface plays an inhibitory role in modulating Ig class switch. This cytokine was shown to enhance cell proliferation of some lymphoma cell lines, while to induce cell death and reduce cell proliferation of other lymphoma cell lines. The pleiotropic biologic activities of this cytokine on different CD30+ lymphoma cell lines may play a pathophysiologic role in Hodgkin's and some non-Hodgkin's lymphomas. [provided by RefSeq]

Other Designations CD153 antigen|CD30 antigen ligand|CD30 ligand|OTTHUMP00000022762

Pathway

- [Cytokine-cytokine receptor interaction](#)

Disease

- [Diabetes Mellitus](#)
- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Kidney Failure](#)

- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Myeloma](#)
- [Occupational Diseases](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)