


Full-Length

CD40 (Human) Recombinant Protein (P01)

Catalog # H00000958-P01

Size 25 ug, 10 ug

Applications



Specification

Product Description	Human CD40 full-length ORF (AAH64518, 1 a.a. - 151 a.a.) recombinant protein with GST-tag at N-terminal.
Sequence	MVRLPLQCVLWGCLLTAVHPEPPTACREKQYLINSQCCSLCQPGQKLVSDCTEFTETECCLPCGE SEFLDTWNRETHFHQHKYCDPNLGLRVQQKGTSETDTICTCEEGWHCTSEACESCVLHRSCSP GFGVKQIDICQPHFPKDRGLNLLM
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	42.35
Interspecies Antigen Sequence	Mouse (57); Rat (52)
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Note

Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — CD40

Entrez GeneID [958](#)

GeneBank Accession# [BC064518](#)

Protein Accession# [AAH64518](#)

Gene Name CD40

Gene Alias Bp50, CDW40, MGC9013, TNFRSF5, p50

Gene Description CD40 molecule, TNF receptor superfamily member 5

Omim ID [109535 606843](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor has been found to be essential in mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand, which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction. The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq]

Other Designations

B cell surface antigen CD40|B cell-associated molecule|CD40 antigen|CD40 antigen (TNF receptor superfamily member 5)|CD40 type II isoform|CD40L receptor|OTTHUMP00000031699|OTTHUMP00000031700|nerve growth factor receptor-related B-lymphocyte activation mol

Pathway

- [Allograft rejection](#)
- [Asthma](#)
- [Autoimmune thyroid disease](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Cytokine-cytokine receptor interaction](#)
- [Primary immunodeficiency](#)
- [Systemic lupus erythematosus](#)
- [Toll-like receptor signaling pathway](#)

Disease

- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Autoimmune Diseases](#)
- [Brain Ischemia](#)
- [Bronchiolitis](#)
- [Calcinosis](#)
- [Cardiovascular Diseases](#)
- [Colitis](#)
- [Coronary Artery Disease](#)
- [Coronary Disease](#)
- [Crohn Disease](#)
- [Diabetes Mellitus](#)
- [Diabetic Angiopathies](#)

- [Disease Progression](#)
- [Drug Eruptions](#)
- [Edema](#)
- [Endometriosis](#)
- [Genetic Predisposition to Disease](#)
- [Graves Disease](#)
- [Hashimoto Disease](#)
- [Hematologic Diseases](#)
- [Hepatitis B](#)
- [Hodgkin Disease](#)
- [Hypertension](#)
- [Infant](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Myeloma](#)
- [Multiple Sclerosis](#)
- [Myasthenia Gravis](#)
- [Occupational Diseases](#)
- [Pemphigus](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)

- [Rupture](#)
- [Stroke](#)
- [Thyroiditis](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Wegener Granulomatosis](#)
- [Werner syndrome](#)