

CD68 rabbit monoclonal antibody

Catalog # H00000968-K Size 100 ug x up to 3

Specification

| | |
|--------------------------------|--|
| Product Description | Rabbit monoclonal antibody raised against a human CD68 peptide using ARM Technology. |
| Immunogen | A synthetic peptide of human CD68 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence. |
| Host | Rabbit |
| Library Construction | Non-fusion antibody library from rabbit spleen (ARM Technology). |
| Expression | Overexpression vector and transfection into 293H cell line. |
| Reactivity | Human |
| Purification | Protein A |
| Isotype | IgG |
| Quality Control Testing | Antibody reactive against human CD68 peptide by ELISA and mammalian transfected lysate by Western Blot. |
| Storage Buffer | In 1x PBS, pH 7.4 |
| Storage Instruction | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing. |
| Deliverable | Up to three rabbit IgG clones of 100 ug each will be delivered to customer. |
| Note | 1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request. |

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — CD68

| | |
|---------------------|--|
| Entrez GeneID | 968 |
| GeneBank Accession# | CD68 |
| Gene Name | CD68 |
| Gene Alias | DKFZp686M18236, GP110, SCARD1 |
| Gene Description | CD68 molecule |
| Omim ID | 153634 |
| Gene Ontology | Hyperlink |
| Gene Summary | <p>This gene encodes a 110-kD transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It is a member of the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family. The protein primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. It is a type I integral membrane protein with a heavily glycosylated extracellular domain and binds to tissue- and organ-specific lectins or selectins. The protein is also a member of the scavenger receptor family. Scavenger receptors typically function to clear cellular debris, promote phagocytosis, and mediate the recruitment and activation of macrophages. Alternative splicing results in multiple transcripts encoding different isoforms. [provided by RefSeq]</p> |
| Other Designations | CD68 antigen OTTHUMP00000135285 macrophage antigen CD68 macrosialin scavenger receptor class D, member 1 |

Pathway

- [Lysosome](#)