

# CDC2 rabbit monoclonal antibody

Catalog # H00000983-K      Size 100 ug x up to 3

## Specification

Product Description	Rabbit monoclonal antibody raised against a human CDC2 peptide using ARM Technology.
Immunogen	A synthetic peptide of human CDC2 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen ( <a href="#">ARM Technology</a> ).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human CDC2 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) <sub>2</sub> , IgG, scFv and different Fc and non-Fc conjugates per customer request.

## Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

## Gene Info — CDC2

Entrez GeneID	<a href="#">983</a>
GeneBank Accession#	<a href="#">CDC2</a>
Gene Name	CDC2
Gene Alias	CDC28A, CDK1, DKFZp686L20222, MGC111195
Gene Description	cell division cycle 2, G1 to S and G2 to M
Omim ID	<a href="#">116940</a>
Gene Ontology	<a href="#">Hyperlink</a>
Gene Summary	The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]
Other Designations	OTTHUMP00000019660 cell cycle controller CDC2 cell division control protein 2 homolog cell division cycle 2 protein cyclin-dependent kinase 1 p34 protein kinase

## Pathway

- [Cell cycle](#)
- [Gap junction](#)
- [p53 signaling pathway](#)

## Disease

- [Alzheimer disease](#)
- [Breast cancer](#)

- [Breast Neoplasms](#)
- [Dementia](#)
- [Genetic Predisposition to Disease](#)
- [Lung Neoplasms](#)
- [Pulmonary Disease](#)