

## CDK7 rabbit monoclonal antibody

Catalog # H00001022-K Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human CDK7 peptide using ARM Technology.
Immunogen	A synthetic peptide of human CDK7 is used for rabbit immunization.  Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen ( <u>ARM Technology</u> ).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	lgG
Quality Control Testing	Antibody reactive against human CDK7 peptide by ELISA and mammalian transfected lysate by We stern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit lgG clones of 100 ug each will be delivered to customer.
Note	<ol> <li>Customer may provide cell or tissue lysate for antibody screening.</li> <li>Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)<sub>2</sub>, lgG, scFv and different Fc and non-Fc conjugates per customer request.</li> </ol>

## **Applications**

Western Blot (Transfected lysate)

Protocol Download



ELISA

Gene Info — CDK7	
Entrez GenelD	1022
GeneBank Accession#	CDK7
Gene Name	CDK7
Gene Alias	CAK1, CDKN7, MO15, STK1, p39MO15
Gene Description	cyclin-dependent kinase 7
Omim ID	601955
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a member of the cyclin-dependent protein kinase (CDK) family. CDK family members are highly similar to the gene products of Saccharomyces cerevisiae cdc 28, and Schizosaccharomyces pombe cdc2, and are known to be important regulators of cell cycl e progression. This protein forms a trimeric complex with cyclin H and MAT1, which functions as a Cdk-activating kinase (CAK). It is an essential component of the transcription factor TFIIH, that is i nvolved in transcription initiation and DNA repair. This protein is thought to serve as a direct link b etween the regulation of transcription and the cell cycle. [provided by RefSeq
Other Designations	39 KDa protein kinase Cdk-activating kinase cell division protein kinase 7 cyclin-dependent kinase 7 (MO15 homolog, Xenopus laevis, cdk-activating kinase) homolog of Xenopus MO15 Cdk-activating kinase kinase subunit of CAK serine/threonine kinase stk1 ser

## Pathway

- Cell cycle
- Nucleotide excision repair

## Disease

- Adenocarcinoma
- Ataxia telangiectasia
- Colonic Neoplasms



- Colorectal Neoplasms
- Esophageal Neoplasms
- Genetic Predisposition to Disease
- Kidney Failure
- Lung Neoplasms
- Multiple Sclerosis
- Ovarian Neoplasms
- Pulmonary Disease
- Rectal Neoplasms
- Urinary Bladder Neoplasms
- Werner syndrome