


Full-Length

CRYAA (Human) Recombinant Protein (P01)

Catalog # H00001409-P01

Size 25 ug, 10 ug

Applications



Specification

Product Description

Human CRYAA full-length ORF (NP_000385.1, 1 a.a. - 173 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence

MDVTIQHPWFKRTLGPFPYPSRLFDQFFGEGLFEYDLLPFLSSTISPYRQSLFRTVLDSGISEVRSD
RDKFVIFLDVKHFSPEDLTVKVQDDFVEIHGKHNERQDDHGYSREFHRRYRLPSNVDQSALSCS
LSADGMLTFCGPKIQTGLDATHAERAIPVSREEKPTSAPSS

Host

Wheat Germ (in vitro)

Theoretical MW (kDa)

46.3

Interspecies Antigen Sequence

Mouse (84); Rat (95)

Preparation Method

[in vitro wheat germ expression system](#)

Purification

Glutathione Sepharose 4 Fast Flow

Quality Control Testing

12.5% SDS-PAGE Stained with Coomassie Blue.

Storage Buffer

50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Note

Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — CRYAA

Entrez GeneID	1409
GeneBank Accession#	NM_000394.2
Protein Accession#	NP_000385.1
Gene Name	CRYAA
Gene Alias	CRYA1, HSPB4
Gene Description	crystallin, alpha A
Omim ID	123580
Gene Ontology	Hyperlink

Gene Summary

Crystallins are separated into two classes: taxon-specific, or enzyme, and ubiquitous. The latter class constitutes the major proteins of vertebrate eye lens and maintains the transparency and refractive index of the lens. Since lens central fiber cells lose their nuclei during development, these crystallins are made and then retained throughout life, making them extremely stable proteins. Mammalian lens crystallins are divided into alpha, beta, and gamma families; beta and gamma crystallins are also considered as a superfamily. Alpha and beta families are further divided into acidic and basic groups. Seven protein regions exist in crystallins: four homologous motifs, a connecting peptide, and N- and C-terminal extensions. Alpha crystallins are composed of two gene products: alpha-A and alpha-B, for acidic and basic, respectively. Alpha crystallins can be induced by heat shock and are members of the small heat shock protein (sHSP also known as the HSP20) family. They act as molecular chaperones although they do not renature proteins and release them in the fashion of a true chaperone; instead they hold them in large soluble aggregates. Post-translational modifications decrease the ability to chaperone. These heterogeneous aggregates consist of 30-40 subunits; the alpha-A and alpha-B subunits have a 3:1 ratio, respectively. Two additional functions of alpha crystallins are an autokinase activity and participation in the intracellular architecture. Alpha-A and alpha-B gene products are differentially expressed; alpha-A is preferentially restricted to the lens and alpha-B is expressed widely in many tissues and organs. Defects in this gene cause autosomal dominant congenital cataract (ADCC). [provided by RefSeq]

Other Designations

crystallin, alpha-1|human alphaA-crystallin (CRYA1)

Disease

- [Cataract](#)
- [Genetic Predisposition to Disease](#)