

# CRYBA1 rabbit monoclonal antibody

Catalog # H00001411-K      Size 100 ug x up to 3

## Specification

<b>Product Description</b>	Rabbit monoclonal antibody raised against a human CRYBA1 peptide using ARM Technology.
<b>Immunogen</b>	A synthetic peptide of human CRYBA1 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
<b>Host</b>	Rabbit
<b>Library Construction</b>	Non-fusion antibody library from rabbit spleen ( <a href="#">ARM Technology</a> ).
<b>Expression</b>	Overexpression vector and transfection into 293H cell line.
<b>Reactivity</b>	Human
<b>Purification</b>	Protein A
<b>Isotype</b>	IgG
<b>Quality Control Testing</b>	Antibody reactive against human CRYBA1 peptide by ELISA and mammalian transfected lysate by Western Blot.
<b>Storage Buffer</b>	In 1x PBS, pH 7.4
<b>Storage Instruction</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Deliverable</b>	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
<b>Note</b>	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) <sub>2</sub> , IgG, scFv and different Fc and non-Fc conjugates per customer request.

## Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

## Gene Info — CRYBA1

Entrez GeneID [1411](#)

GeneBank Accession# [CRYBA1](#)

Gene Name CRYBA1

Gene Alias CRYB1

Gene Description crystallin, beta A1

Omim ID [123610 600881](#)

Gene Ontology [Hyperlink](#)

**Gene Summary**

Crystallins are separated into two classes: taxon-specific, or enzyme, and ubiquitous. The latter class constitutes the major proteins of vertebrate eye lens and maintains the transparency and refractive index of the lens. Since lens central fiber cells lose their nuclei during development, these crystallins are made and then retained throughout life, making them extremely stable proteins. Mammalian lens crystallins are divided into alpha, beta, and gamma families; beta and gamma crystallins are also considered as a superfamily. Alpha and beta families are further divided into acidic and basic groups. Seven protein regions exist in crystallins: four homologous motifs, a connecting peptide, and N- and C-terminal extensions. Beta-crystallins, the most heterogeneous, differ by the presence of the C-terminal extension (present in the basic group, none in the acidic group). Beta-crystallins form aggregates of different sizes and are able to self-associate to form dimers or to form heterodimers with other beta-crystallins. This gene, a beta acidic group member, encodes two proteins (crystallin, beta A3 and crystallin, beta A1) from a single mRNA, the latter protein is 17 aa shorter than crystallin, beta A3 and is generated by use of an alternate translation initiation site. Deletion of exons 3 and 4 causes the autosomal dominant disease 'zonular cataract with sutural opacities'. [provided by RefSeq]

**Other Designations** crystallin, beta A3|eye lens structural protein