

DAB2 rabbit monoclonal antibody

Catalog # H00001601-K Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human DAB2 peptide using ARM Technology.
Immunogen	A synthetic peptide of human DAB2 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human DAB2 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — DAB2

Entrez GeneID [1601](#)

GeneBank Accession# [DAB2](#)

Gene Name DAB2

Gene Alias DOC-2, DOC2, FLJ26626

Gene Description disabled homolog 2, mitogen-responsive phosphoprotein (Drosophila)

Omim ID [601236](#)

Gene Ontology [Hyperlink](#)

Gene Summary

DAB2 mRNA is expressed in normal ovarian epithelial cells but is down-regulated or absent from ovarian carcinoma cell lines. The 770-amino acid predicted protein has an overall 83% identity with the mouse p96 protein, a putative mitogen-responsive phosphoprotein; homology is strongest in the amino-terminal end of the protein in a region corresponding to the phosphotyrosine interaction domain. The down-regulation of DAB2 may play an important role in ovarian carcinogenesis. This gene was initially named DOC2 (for Differentially expressed in Ovarian Cancer) and is distinct from the DOC2A and DOC2B genes (for double C2-like domains, alpha and beta). [provided by RefSeq]

Other Designations disabled homolog 2|mitogen-responsive phosphoprotein

Pathway

- [Endocytosis](#)

Disease

- [Alzheimer disease](#)
- [Birth Weight](#)
- [Cerebral Amyloid Angiopathy](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)

- [Glioma](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Neuroblastoma](#)
- [Ovarian Neoplasms](#)
- [Tobacco Use Disorder](#)