


MaxPab®

DBT purified MaxPab rabbit polyclonal antibody (D01P)

Catalog # H00001629-D01P


Size 100 ug

Applications




Western Blot (Tissue lysate)

DBT MaxPab rabbit polyclonal antibody. Western Blot analysis of DBT expression in human kidney.



Western Blot (Cell lysate)

DBT MaxPab rabbit polyclonal antibody. Western Blot analysis of DBT expression in A-431.



Western Blot (Transfected lysate)

Western Blot analysis of DBT expression in transfected 293T cell line ([H00001629-T02](#)) by DBT MaxPab polyclonal antibody.

Lane 1: DBT transfected lysate(53.50 KDa).

Lane 2: Non-transfected lysate.

Specification

Product Description

Rabbit polyclonal antibody raised against a full-length human DBT protein.

Immunogen	DBT (AAH16675.1, 1 a.a. ~ 482 a.a) full-length human protein.
Sequence	MAAVRMLRTWSRNAGKLCVRYFQTCGNVHVLKPNYVCFEGYPSFKYSHPHHFLKTTAALRGQV VQFKLSDIGEGIREVTVKEWYVKEGDTV SQFDSICEVQSDKASVTITSRYDGVKKLYNLDDIAYV GKPLVDIETEALKDSEEDVVETPAVSHDEHHTQEIKGKRTLATPAVRRLAMENNIKLSEVVGSGK DGRILKEDILNYLEKQTGAILPPSPKVEIMPPPKPKDMTPILVSKPPVFTGKDKTEPIKGFQKAM VKTMSAALKIPHFGYCDEIDLTELVLKREELKPIAFARGIKLSFMPFFLKAASLGLLQFPILNASVDE NCQNITYKASHNIGIAMDTEQGLVPNVKNVQICSIFDIATELNRLQKLGSVQQLSTTDLTG GTFTLSNI GSIGGTFAKPVIMPPEVAIGALGSIKAIPRFNQKGEVYKAQIMNVSW SADHRVIDGATMSRFSNLW KSYLENPAFMLLDLK
Host	Rabbit
Reactivity	Human
Interspecies Antigen Sequence	Mouse (88); Rat (88)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Tissue lysate)

DBT MaxPab rabbit polyclonal antibody. Western Blot analysis of DBT expression in human kidney.

[Protocol Download](#)

- Western Blot (Cell lysate)

DBT MaxPab rabbit polyclonal antibody. Western Blot analysis of DBT expression in A-431.

[Protocol Download](#)

- Western Blot (Transfected lysate)

Western Blot analysis of DBT expression in transfected 293T cell line ([H00001629-T02](#)) by DBT MaxPab polyclonal antibody.

Lane 1: DBT transfected lysate(53.50 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

Gene Info — DBT

Entrez GeneID	1629
GeneBank Accession#	BC016675.1
Protein Accession#	AAH16675.1
Gene Name	DBT
Gene Alias	BCATE2, E2, E2B, MGC9061
Gene Description	dihydrolipoamide branched chain transacylase E2
Omim ID	248600 248610
Gene Ontology	Hyperlink
Gene Summary	The branched-chain alpha-keto acid dehydrogenase complex (BCKD) is an inner-mitochondrial enzyme complex involved in the breakdown of the branched-chain amino acids isoleucine, leucine, and valine. The BCKD complex is thought to be composed of a core of 24 transacylase (E2) subunits, and associated decarboxylase (E1), dehydrogenase (E3), and regulatory subunits. This gene encodes the transacylase (E2) subunit. Mutations in this gene result in maple syrup urine disease, type 2. Alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq]
Other Designations	BCKAD E2 subunit E2 component of branched chain alpha-keto acid dehydrogenase complex O TTHUMP00000012596 branched chain acyltransferase, E2 component dihydrolipoamide branched chain transacylase dihydrolipoyl transacylase dihydrolipoyllysine-residue (2-me

Pathway

- [Metabolic pathways](#)
- [Valine](#)