DDX5 rabbit monoclonal antibody

Catalog # H00001655-K

ocification

Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human DDX5 peptide using ARM Technology.
Immunogen	A synthetic peptide of human DDX5 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
lsotype	lgG
Quality Control Testing	Antibody reactive against human DDX5 peptide by ELISA and mammalian transfected lysate by We stern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, lgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

• Western Blot (Transfected lysate)

Protocol Download



• ELISA

Gene Info — DDX5	
Entrez GenelD	<u>1655</u>
GeneBank Accession#	DDX5
Gene Name	DDX5
Gene Alias	DKFZp434E109, DKFZp686J01190, G17P1, HLR1, HUMP68, p68
Gene Description	DEAD (Asp-Glu-Ala-Asp) box polypeptide 5
Omim ID	180630
Gene Ontology	Hyperlink
Gene Summary	DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure, such as translation initiation, nuclear and mitochondrial splicing, and ribosom e and spliceosome assembly. Based on their distribution patterns, some members of this family a re believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein, which is a RNA-dependent ATPase, and also a prolifera tion-associated nuclear antigen, specifically reacting with the simian virus 40 tumor antigen. This gene consists of 13 exons, and alternatively spliced transcripts containing several intron sequenc es have been detected, but no isoforms encoded by these transcripts have been identified. [provi ded by RefSeq
Other Designations	ATP-dependent RNA helicase DDX5 DEAD box-5 DEAD/H (Asp-Glu-Ala-Asp/His) box polypepti de 5 (RNA helicase, 68kD)

Disease

- Disease Progression
- Genetic Predisposition to Disease
- Hepatitis C
- Kidney Failure
- Liver Cirrhosis