

DNAxPAb



DUSP6 DNAxPab

Catalog # H00001848-W01P Size 200 ug

Specification	
Product Description	Rabbit polyclonal antibody raised against a full-length human DUSP6 DNA using DNAx™ Immune te chnology.
Technology	DNAx™ Immune
Immunogen	Full-length human DNA
Sequence	MIDTLRPVPFASEMAISKTVAWLNEQLELGNERLLLMDCRPQELYESSHIESAINVAIPGIMLRRLQ KGNLPVRALFTRGEDRDRFTRRCGTDTVVLYDESSSDWNENTGGESVLGLLLKKLKDEGCRAFY LEGGFSKFQAEFSLHCETNLDGSCSSSSPPLPVLGLGGGLRISSDSSSDIESDLDRDPNSATDSD GSPLSNSQPSFPVEILPFLYLGCAKDSTNLDVLEEFGIKYILNVTPNLPNLFENAGEFKYKQIPISDH WSQNLSQFFPEAISFIDEARGKNCGVLVHCLAGISRSVTVTVAYLMQKLNLSMNDAYDIVKMKKSN ISPNFNFMGQLLDFERTLGLSSPCDNRVPAQQLYFTTPSNQNVYQVDSLQST
Host	Rabbit
Reactivity	Human
Purification	Protein A
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

Western Blot (Transfected lysate)

Protocol Download

• Immunofluorescence (Transfected cell)

• Flow Cytometry (Transfected cell)

Gene Info — DUSP6	
Entrez GenelD	<u>1848</u>
GeneBank Accession#	<u>NM_001946.2</u>
Protein Accession#	<u>NP_001937.2</u>
Gene Name	DUSP6
Gene Alias	MKP3, PYST1
Gene Description	dual specificity phosphatase 6
Omim ID	<u>602748</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a member of the dual specificity protein phosphatase subfam ily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoser ine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-ac tivated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated w ith cellular proliferation and differentiation. Different members of the family of dual specificity phos phatases show distinct substrate specificities for various MAP kinases, different tissue distributio n and subcellular localization, and different modes of inducibility of their expression by extracellula r stimuli. This gene product inactivates ERK2, is expressed in a variety of tissues with the highest levels in heart and pancreas, and unlike most other members of this family, is localized in the cyto plasm. Two transcript variants encoding different isoforms have been found for this gene. [provide d by RefSeq

Pathway

• MAPK signaling pathway

Disease

- Bipolar Disorder
- <u>Cardiovascular Diseases</u>

😵 Abnova

- Diabetes Mellitus
- Edema
- Genetic Predisposition to Disease
- Kidney Failure