. .

E2F3 rabbit monoclonal antibody

Catalog # H00001871-K Size 100 ug x up to 3

Specification	
Product Description	Rabbit monoclonal antibody raised against a human E2F3 peptide using ARM Technology.
Immunogen	A synthetic peptide of human E2F3 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
lsotype	lgG
Quality Control Testing	Antibody reactive against human E2F3 peptide by ELISA and mammalian transfected lysate by Wes tern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

• Western Blot (Transfected lysate)

Protocol Download



• ELISA

Gene Info — E2F3	
Entrez GenelD	<u>1871</u>
GeneBank Accession#	<u>E2F3</u>
Gene Name	E2F3
Gene Alias	DKFZp686C18211, E2F-3, KIAA0075, MGC104598
Gene Description	E2F transcription factor 3
Omim ID	<u>600427</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain s everal evolutionally conserved domains found in most members of the family. These domains incl ude a DNA binding domain, a dimerization domain which determines interaction with the different iation regulated transcription factor proteins (DP), a transactivation domain enriched in acidic ami no acids, and a tumor suppressor protein association domain which is embedded within the trans activation domain. This protein and another 2 members, E2F1 and E2F2, have an additional cycli n binding domain. This protein binds specifically to retinoblastoma protein pRB in a cell-cycle dep endent manner. [provided by RefSeq
Other Designations	OTTHUMP00000018012

Pathway

- Bladder cancer
- <u>Cell cycle</u>
- Chronic myeloid leukemia
- <u>Glioma</u>
- Melanoma
- Non-small cell lung cancer
- Pancreatic cancer



- Pathways in cancer
- Prostate cancer
- Small cell lung cancer

Disease

- Genetic Predisposition to Disease
- Ovarian Neoplasms