


Full-Length

ENO1 (Human) Recombinant Protein (P02)

Catalog # H00002023-P02

Size 25 ug, 10 ug

Applications



Specification

Product Description

Human ENO1 full-length ORF (AAH22545, 1 a.a. - 434 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence

MSILKIHAREIFDSRGNPTVEVDLFTSKGLFRAAVPSGASTGIYEALERDNDKTRYMGKGVSKAV
EHINKTIAPALVSKKLVTEQEKIDKLMIMDGTENKSKFGANILGVSLAVCKAGAVEKGVPLYRHI
ADLAGNSEVILPVPAFNVINGGSHAGNKLAMQEFMILPVGAANFREAMRIGAEVYHNLKNVIKEY
GKDATNVGDEGGFAPNILENKEGLELLKTAIGKAGYTDKVVIGMDVAASEFFRSGKYDLDFKSPD
DPSRYISPDQLADLYKSFIDYPVVSIEDPFDQDDWGAWQKFTASAGIQVVGDDLTVTNPKRIKA
VNEKSCNCLLLKVNQIGSVTESLQACKLAQANGWGMVSHRSGETEDTFIADLVVGLCTGQIKTG
APCRSERLAKYNQLLRIEEEELGSKAKFAGRNFRNPLAK

Host

Wheat Germ (in vitro)

Theoretical MW (kDa)

73.48

Preparation Method

[in vitro wheat germ expression system](#)

Purification

Glutathione Sepharose 4 Fast Flow

Quality Control Testing

12.5% SDS-PAGE Stained with Coomassie Blue.

Storage Buffer

50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction

Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Note

Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — ENO1

Entrez GeneID [2023](#)

GeneBank Accession# [BC022545](#)

Protein Accession# [AAH22545](#)

Gene Name ENO1

Gene Alias ENO1L1, MBP-1, MPB1, NNE, PPH

Gene Description enolase 1, (alpha)

Omim ID [172430](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene encodes one of three enolase isoenzymes found in mammals; it encodes alpha-enolase, a homodimeric soluble enzyme, and also encodes a shorter monomeric structural lens protein, tau-crystallin. The two proteins are made from the same message. The full length protein, the isoenzyme, is found in the cytoplasm. The shorter protein is produced from an alternative translation start, is localized to the nucleus, and has been found to bind to an element in the c-myc promoter. A pseudogene has been identified that is located on the other arm of the same chromosome. [provided by RefSeq]

Other Designations 2-phospho-D-glycerate hydro-lyase|MYC promoter-binding protein 1|OTTHUMP00000001706|alpha enolase like 1|enolase 1|non-neural enolase|phosphopyruvate hydratase|tau-crystallin

Pathway

- [Biosynthesis of alkaloids derived from histidine and purine](#)
- [Biosynthesis of alkaloids derived from ornithine](#)
- [Biosynthesis of alkaloids derived from shikimate pathway](#)
- [Biosynthesis of alkaloids derived from terpenoid and polyketide](#)
- [Biosynthesis of phenylpropanoids](#)
- [Biosynthesis of plant hormones](#)
- [Biosynthesis of terpenoids and steroids](#)
- [Glycolysis / Gluconeogenesis](#)
- [Metabolic pathways](#)
- [RNA degradation](#)

Disease

- [Myocardial Infarction](#)