
EPHB6 (Human) Recombinant Protein (Q01)

Catalog # H00002051-Q01 Size 25 ug, 10 ug

Applications



| Specification | |
|----------------------------------|---|
| Product Description | Human EPHB6 partial ORF (NP_004436, 23 a.a 122 a.a.) recombinant protein with GST-tag at N- terminal. |
| Sequence | DTTGETSEIGWLTYPPGGWDEVSVLDDQRRLTRTFEACHVAGAPPGTGQDNWLQTHFVERRGA QRAHIRLHFSVRACSSLGVSGGTCRETFTLYYRQAEE |
| Host | Wheat Germ (in vitro) |
| Theoretical MW (kDa) | 36.63 |
| Interspecies Antigen Sequence | Mouse (97); Rat (97) |
| Preparation Method | in vitro wheat germ expression system |
| Purification | Glutathione Sepharose 4 Fast Flow |
| Quality Control Testing | 12.5% SDS-PAGE Stained with Coomassie Blue. |
| Storage Buffer | 50 mM Tris-HCI, 10 mM reduced Glutathione, pH=8.0 in the elution buffer. |
| Storage Instruction | Store at -80°C. Aliquot to avoid repeated freezing and thawing. |
| Note | Best use within three months from the date of receipt of this protein. |



Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

| Gene Info — EPHB6 | |
|---------------------|---|
| Entrez GenelD | <u>2051</u> |
| GeneBank Accession# | <u>NM_004445</u> |
| Protein Accession# | <u>NP_004436</u> |
| Gene Name | EPHB6 |
| Gene Alias | HEP, MGC129910, MGC129911 |
| Gene Description | EPH receptor B6 |
| Omim ID | <u>602757</u> |
| Gene Ontology | Hyperlink |
| Gene Summary | Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, par ticularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosp hatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The E ph family of receptors are divided into 2 groups based on the similarity of their extracellular domai n sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptor smake up the largest subgroup of the receptor tyrosine kinase (RTK) family. The ephrin receptor encode d by this gene lacks the kinase activity of most receptor tyrosine kinases and binds to ephrin-B lig ands. [provided by RefSeq |
| Other Designations | ephrin receptor EphB6 |

Pathway

• Axon guidance