


MaxPab®

FGFR1 purified MaxPab rabbit polyclonal antibody (D01P)

Catalog # H00002260-D01P Size 100 ug

Applications




Western Blot (Transfected lysate)

Western Blot analysis of FGFR1 expression in transfected 293T cell line ([H00002260-T03](#)) by FGFR1 MaxPab polyclonal antibody.

Lane 1: FGFR1 transfected lysate(81.90 KDa).

Lane 2: Non-transfected lysate.



In situ Proximity Ligation Assay (Cell)

Proximity Ligation Analysis of protein-protein interactions between FGFR1 and CTNNB1. HeLa cells were stained with anti-FGFR1 rabbit purified polyclonal 1:1200 and anti-CTNNB1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

Specification

| | |
|---------------------|--|
| Product Description | Rabbit polyclonal antibody raised against a full-length human FGFR1 protein. |
|---------------------|--|

| | |
|-----------|--|
| Immunogen | FGFR1 (NP_075594.1, 1 a.a. ~ 731 a.a) full-length human protein. |
|-----------|--|

| | |
|--------------------------------|---|
| Sequence | MWSWKCLLFWAVLTATLCTARPSPTLPEQDALPSSSEDDDDDDSSSEEKETDNTKPNPVAPY WTSPEKMEKKLHAVPAAKTVFKCPSSGTPNPTLRWLNGKEFPDHRIGGYKVRYATWSIIMDS V/PSDKGNYTCIVENEYGSINHTYQLDVVERSPHRPILQAGLPANKTVALGSNVEFMCKVYSDPQ PHIQWLKHIEVNGSKIGPDNLPYVQILKTAGVNNDKEMEVVLHRLNVSFEDAGEYTCLAGNSIGLSH HSAWLTVLEALEERPAVMTSPLYLEIIYCTGAFLISCMGVSVIVYMKSGTKKSDFHQSMAVHKLA KSIPLRRQVTVSADSSASMNSGVLLVRPSRLSSSGTPMLAGVSEYELPEDPRWELPRDRLVLGK PLGEGCFGQVVLAEAGLDKDKPNRVTKAVKMLKSDATEKDLSDLISEMEMMKMIGHKNIINLL GACTQDGPLYVIVEYASKGNLREYLQARRPGLEYCYNPSHNPEEQLSSKDLVSCAYQVARGME YLASKKCIHRDLAARNVLVTEDNVMKIADFGLARDIHHIDYYKKTTNGRLPVKWMMAPEALFDRIYTH QSDVWSFGVLLWEITLGGSPYGPVVEELFKLLKEGHRMDKPSNCTNELYMMMRDCWHAVPS QRPTFKQLVEDLDRIVALTSNQEYLDLSMP LDQYSPSFPDTRSSTCSSGEDSVFSHEPLPEEPCL PRHPAQLANGGLKRR |
| Host | Rabbit |
| Reactivity | Human |
| Quality Control Testing | Antibody reactive against mammalian transfected lysate. |
| Storage Buffer | In 1x PBS, pH 7.4 |
| Storage Instruction | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing. |

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of FGFR1 expression in transfected 293T cell line ([H00002260-T03](#)) by FGFR1 MaxPab polyclonal antibody.

Lane 1: FGFR1 transfected lysate(81.90 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

- In situ Proximity Ligation Assay (Cell)

Proximity Ligation Analysis of protein-protein interactions between FGFR1 and CTNNB1. HeLa cells were stained with anti-FGFR1 rabbit purified polyclonal 1:1200 and anti-CTNNB1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

Gene Info — FGFR1

| | |
|---------------------|-----------------------------|
| Entrez GenelD | 2260 |
| GeneBank Accession# | NM_023106.2 |
| Protein Accession# | NP_075594.1 |

| | |
|--------------------|---|
| Gene Name | FGFR1 |
| Gene Alias | BFGFR, CD331, CEK, FGFBP, FLG, FLJ99988, FLT2, HBGFR, KAL2, N-SAM |
| Gene Description | fibroblast growth factor receptor 1 |
| Omim ID | 101600 123150 136350 147950 |
| Gene Ontology | Hyperlink |
| Gene Summary | The protein encoded by this gene is a member of the fibroblast growth factor receptor (FGFR) family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member binds both acidic and basic fibroblast growth factors and is involved in limb induction. Mutations in this gene have been associated with Pfeiffer syndrome, Jackson-Weiss syndrome, Antley-Bixler syndrome, osteoglophonic dysplasia, and autosomal dominant Kallmann syndrome 2. Chromosomal aberrations involving this gene are associated with stem cell myeloproliferative disorder and stem cell leukemia lymphoma syndrome. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized. [provided by RefSeq] |
| Other Designations | FMS-like tyrosine kinase 2 OTTHUMP00000190874 OTTHUMP00000190878 OTTHUMP00000190879 OTTHUMP00000190881 basic fibroblast growth factor receptor 1 fms-related tyrosine kinase 2 fms-related tyrosine kinase-2 heparin-binding growth factor receptor hydroxyaryl |

Pathway

- [Adherens junction](#)
- [MAPK signaling pathway](#)
- [Melanoma](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Regulation of actin cytoskeleton](#)

Disease

- [Abnormalities](#)

- [Acrocephalosyndactyla](#)
- [Alzheimer disease](#)
- [Amenorrhea](#)
- [Anodontia](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)
- [Cardiovascular Diseases](#)
- [Chromosome Aberrations](#)
- [Chromosome Disorders](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Craniofacial Dysostosis](#)
- [Craniosynostoses](#)
- [Diabetes Complications](#)
- [Fractures](#)
- [Genetic Diseases](#)
- [Genetic Predisposition to Disease](#)
- [Hypersensitivity](#)
- [Hypogonadism](#)
- [Kallmann Syndrome](#)
- [Metabolic Syndrome X](#)
- [Neoplasms](#)
- [Obesity](#)
- [Osteoporosis](#)
- [Ovarian Failure](#)

- [Polycystic Ovary Syndrome](#)
- [Puberty](#)
- [Schizophrenia](#)
- [Sleep Apnea](#)
- [Thrombophilia](#)
- [Thyroid Neoplasms](#)
- [Tobacco Use Disorder](#)
- [Tooth Abnormalities](#)