


FGFR2 monoclonal antibody (M04), clone 3F4

Catalog # H00002263-M04

Size 100 ug

Applications



Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged FGFR2 is approximately 0.1ng/ml as a capture antibody.

Specification

Product Description	Mouse monoclonal antibody raised against a partial recombinant FGFR2.
Immunogen	FGFR2 (AAH39243, 621 a.a. ~ 723 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	GHRMDKPANCTNELYMMMRDCWHAVPSQRPTFKQLVEDLDRILTLTTNEEYLDLSQPLEQYSPS YPDTRSSCSSGDDSVFSPDPMPYEPCLPQYPHINGSVKT
Host	Mouse
Reactivity	Human
Interspecies Antigen Sequence	Mouse (98); Rat (97)
Isotype	IgG2b Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged FGFR2 is approximately 0.1ng/ml as a capture antibody.

[Protocol Download](#)

- ELISA

Gene Info — FGFR2

Entrez GeneID	2263
GeneBank Accession#	BC039243
Protein Accession#	AAH39243
Gene Name	FGFR2
Gene Alias	BEK, BFR-1, CD332, CEK3, CFD1, ECT1, FLJ98662, JWS, K-SAM, KGFR, TK14, TK25
Gene Description	fibroblast growth factor receptor 2
Omim ID	101200 101400 101600 123150 123500 123790 137215 149730 176943 207410
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the fibroblast growth factor receptor family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member is a high-affinity receptor for acidic, basic and/or keratinocyte growth factor, depending on the isoform. Mutations in this gene are associated with Crouzon syndrome, Pfeiffer syndrome, Craniosynostosis, Apert syndrome, Jackson-Weiss syndrome, Beare-Stevenson cutis gyrata syndrome, Saethre-Chotzen syndrome, and syndromic craniosynostosis. Multiple alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq]
Other Designations	BEK fibroblast growth factor receptor FGF receptor OTTHUMP00000020621 OTTHUMP00000020629 bacteria-expressed kinase hydroxyaryl-protein kinase keratinocyte growth factor receptor protein tyrosine kinase, receptor like 14 soluble FGFR4 variant 4

Pathway

- [Endocytosis](#)
- [MAPK signaling pathway](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Regulation of actin cytoskeleton](#)

Disease

- [Acrocephalosyndactylia](#)
- [Adenocarcinoma](#)
- [Alzheimer disease](#)
- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Craniofacial Dysostosis](#)
- [Craniosynostoses](#)
- [Cystadenocarcinoma](#)
- [Depressive Disorder](#)
- [Diabetes Complications](#)
- [Disease Susceptibility](#)

- [Endometrial Neoplasms](#)
- [Endometriosis](#)
- [Genetic Predisposition to Disease](#)
- [Hypersensitivity](#)
- [Hypospadias](#)
- [Lymphatic Metastasis](#)
- [Malignant melanoma](#)
- [Melanoma](#)
- [Metabolic Syndrome X](#)
- [Neoplasm Invasiveness](#)
- [Neoplasms](#)
- [Osteoporosis](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Pancreatic cancer](#)
- [Pancreatic Neoplasms](#)
- [Schizophrenia](#)
- [Skin Neoplasms](#)
- [Sleep Apnea](#)
- [Tobacco Use Disorder](#)