FKBP4 (Human) Recombinant Protein (Q01)

Catalog # H00002288-Q01 Size 25 ug, 10 ug

Applications



Specification	
Product Description	Human FKBP4 partial ORF (AAH07924, 301 a.a 410 a.a.) recombinant protein with GST-tag at N -terminal.
Sequence	EYESSFSNEEAQKAQALRLASHLNLAMCHLKLQAFSAAIESCNKALELDSNNEKGLFRRGEAHLA VNDFELARADFQKVLQLYPNNKAAKTQLAVCQQRIRRQLAREKKL
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	37.84
Interspecies Antigen Sequence	Mouse (94); Rat (94)
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCI, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.



Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — FKBP4	
Entrez GenelD	2288
GeneBank Accession#	<u>BC007924</u>
Protein Accession#	<u>AAH07924</u>
Gene Name	FKBP4
Gene Alias	FKBP52, FKBP59, HBI, Hsp56, PPlase, p52
Gene Description	FK506 binding protein 4, 59kDa
Omim ID	<u>600611</u>
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a member of the immunophilin protein family, which play a rol e in immunoregulation and basic cellular processes involving protein folding and trafficking. This e ncoded protein is a cis-trans prolyl isomerase that binds to the immunosuppressants FK506 and r apamycin. It has high structural and functional similarity to FK506-binding protein 1A (FKBP1A), b ut unlike FKBP1A, this protein does not have immunosuppressant activity when complexed with F K506. It interacts with interferon regulatory factor-4 and plays an important role in immunoregulato ry gene expression in B and T lymphocytes. This encoded protein is known to associate with phyt anoyl-CoA alpha-hydroxylase. It can also associate with two heat shock proteins (hsp90 and hsp7 0) and thus may play a role in the intracellular trafficking of hetero-oligomeric forms of the steroid h ormone receptors. This protein correlates strongly with adeno-associated virus type 2 vectors (AA V) resulting in a significant increase in AAV-mediated transgene expression in human cell lines. T hus this encoded protein is thought to have important implications for the optimal use of AAV vect ors in human gene therapy. The human genome contains several non-transcribed pseudogenes si milar to this gene. [provided by RefSeq
Other Designations	52 kD FK506 binding protein FK506 binding protein 4 (59kD) FK506 binding protein 52 FK506-b inding protein 4 (59kD) HSP binding immunophilin T-cell FK506-binding protein, 59kD p59 protei n peptidylprolyl cis-trans isomerase rotamase



Disease

- Asthma
- Genetic Predisposition to Disease
- Glaucoma
- <u>Hypospadias</u>
- <u>Syndrome</u>