


FLT1 (Human) Recombinant Protein (Q01)

Catalog # H00002321-Q01

Size 25 ug, 10 ug

Applications



Specification

Product Description	Human FLT1 partial ORF (AAH39007, 581 a.a. - 687 a.a.) recombinant protein with GST-tag at N-terminal.
Sequence	KFLYRDVTWILLRTVNNRTMHYSISKQKMAITKEHSITLNLTIMNVSLQDSGTACRARNVYTGEEILQKKEITIRGEHCNKKAVFSRISKFKSTRNDCTTQSNVKH
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	37.51
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Quality Control Testing	12.5% SDS-PAGE Stained with Coomassie Blue.
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — FLT1

Entrez GeneID	2321
GeneBank Accession#	BC039007
Protein Accession#	AAH39007
Gene Name	FLT1
Gene Alias	FLT, VEGFR1
Gene Description	fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor)
Omim ID	165070
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a member of the vascular endothelial growth factor receptor (VEGFR) family. VEGFR family members are receptor tyrosine kinases (RTKs) which contain an extracellular ligand-binding region with seven immunoglobulin (Ig)-like domains, a transmembrane segment, and a tyrosine kinase (TK) domain within the cytoplasmic domain. This protein binds to VEGFR-A, VEGFR-B and placental growth factor and plays an important role in angiogenesis and vasculogenesis. Expression of this receptor is found in vascular endothelial cells, placental trophoblast cells and peripheral blood monocytes. Multiple transcript variants encoding different isoforms have been found for this gene. Isoforms include a full-length transmembrane receptor isoform and shortened, soluble isoforms. The soluble isoforms are associated with the onset of pre-eclampsia
Other Designations	fms-related tyrosine kinase 1 soluble VEGF receptor 1-14 soluble VEGFR1 variant 2 soluble VEGFR1 variant 21 vascular endothelial growth factor/vascular permeability factor receptor

Pathway

- [Cytokine-cytokine receptor interaction](#)

- [Endocytosis](#)
- [Focal adhesion](#)

Disease

- [Abortion](#)
- [Adenocarcinoma](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)
- [Cardiovascular Diseases](#)
- [Cell Transformation](#)
- [Chorioamnionitis](#)
- [Colorectal Neoplasms](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Fetal Growth Retardation](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Hypercholesterolemia](#)
- [Hypersensitivity](#)
- [Inflammation](#)
- [Kidney Failure](#)
- [Lymphoma](#)
- [Malaria](#)
- [Melanoma](#)

- [Neovascularization](#)
- [Obstetric Labor](#)
- [Placenta Diseases](#)
- [Pre-Eclampsia](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Sarcoidosis](#)
- [Scleroderma](#)
- [Skin Neoplasms](#)
- [Vaginosis](#)