CCR10 rabbit monoclonal antibody

Catalog # H00002826-K

Specification

Size 100 ug x up to 3

opecification	
Product Description	Rabbit monoclonal antibody raised against a human CCR10 peptide using ARM Technology.
Immunogen	A synthetic peptide of human CCR10 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
lsotype	lgG
Quality Control Testing	Antibody reactive against human CCR10 peptide by ELISA and mammalian transfected lysate by W estern Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	 Customer may provide cell or tissue lysate for antibody screening. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering in cluding F(ab)₂, IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

• Western Blot (Transfected lysate)

Protocol Download

• ELISA

Gene Info — CCR10	
Entrez GenelD	2826
GeneBank Accession#	CCR10
Gene Name	CCR10
Gene Alias	GPR2
Gene Description	chemokine (C-C motif) receptor 10
Omim ID	<u>600240</u>
Gene Ontology	Hyperlink
Gene Summary	Hyperlink Chemokines are a group of small (approximately 8 to 14 kD), mostly basic, structurally related mo lecules that regulate cell trafficking of various types of leukocytes through interactions with a subse t of 7-transmembrane, G protein-coupled receptors. Chemokines also play fundamental roles in th e development, homeostasis, and function of the immune system, and they have effects on cells of the central nervous system as well as on endothelial cells involved in angiogenesis or angiostasis. Chemokines are divided into 2 major subfamilies, CXC and CC, based on the arrangement of th e first 2 of the 4 conserved cysteine residues; the 2 cysteines are separated by a single amino aci d in CXC chemokines and are adjacent in CC chemokines. CCR10 is the receptor for CCL27 (S CYA27; MIM 604833); CCR10-CCL27 interactions are involved in T cell-mediated skin inflammat ion (Homey et al., 2002 [PubMed 11821900]).[supplied by OMIM

Pathway

- Chemokine signaling pathway
- Cytokine-cytokine receptor interaction