


HCK (Human) IP-WB Antibody Pair

Catalog # H00003055-PW1 Size 1 Set

Applications



Immunoprecipitation of HCK transfected lysate using rabbit polyclonal anti-HCK and Protein A Magnetic Bead (<u>U0007</u>), and immunoblotted with mouse purified polyclonal anti-HCK.

Specification	
Product Description	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.
Reactivity	Human
Quality Control Testing	Immunoprecipitation-Western Blot (IP-WB) Immunoprecipitation of HCK transfected lysate using rabbit polyclonal anti-HCK and Protein A Magn etic Bead (U0007), and immunoblotted with mouse purified polyclonal anti-HCK.
Supplied Product	Antibody pair set content: 1. Antibody pair for IP: rabbit polyclonal anti-HCK (300 ul) 2. Antibody pair for WB: mouse purified polyclonal anti-HCK (50 ug)
Storage Instruction	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze tha w cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

Immunoprecipitation-Western Blot

Protocol Download



Gene Info — HCK	
Entrez GenelD	<u>3055</u>
Gene Name	HCK
Gene Alias	JTK9
Gene Description	hemopoietic cell kinase
Omim ID	<u>142370</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a protein-tyrosine kinase that is predominantly expressed in hemopoietic cell types. The encoded protein may help couple the Fc receptor to the activation of the respiratory burst. In addition, it may play a role in neutrophil migration and in the degranulation of neutrophils. Alternate translation initiation site usage, including a non-AUG (CUG) codon, result in the production of two different isoforms, that have different subcellular localization. [provided by RefSeq
Other Designations	tyrosine protein kinase HCK

Pathway

- Chemokine signaling pathway
- Fc gamma R-mediated phagocytosis

Disease

- HIV Infections
- Pulmonary Disease