


HLA-DMB monoclonal antibody (M06A), clone 5G11

Catalog # H00003109-M06A

Size 200 uL

Applications




Western Blot (Transfected lysate)

Western Blot analysis of HLA-DMB expression in transfected 293T cell line by HLA-DMB monoclonal antibody (M06A), clone 5G11.

Lane 1: HLA-DMB transfected lysate(28.9 KDa).

Lane 2: Non-transfected lysate.



Western Blot detection against Immunogen (37.62 KDa) .

Specification

Product Description	Mouse monoclonal antibody raised against a partial recombinant HLA-DMB.
Immunogen	HLA-DMB (NP_002109, 22 a.a. ~ 129 a.a) partial recombinant protein with GST tag. MW of the GS T tag alone is 26 KDa.
Sequence	VAHVESTCLDDAGTPKDFTYCISFNKDLLTCWDPEENKMAPCEFGVLNSLANVLSQHLNQKDT LMQRLRNLQNCATHTQPFWGSLTNRTRPPSVQVAKTTPFNTRE
Host	Mouse
Reactivity	Human
Isotype	IgG2b Kappa

Quality Control Testing

Antibody Reactive Against Recombinant Protein.
Western Blot detection against Immunogen (37.62 KDa) .

Storage Buffer

In ascites fluid

Storage Instruction

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Transfected lysate)

Western Blot analysis of HLA-DMB expression in transfected 293T cell line by HLA-DMB monoclonal antibody (M06A), clone 5G11.

Lane 1: HLA-DMB transfected lysate(28.9 KDa).

Lane 2: Non-transfected lysate.

[Protocol Download](#)

- Western Blot (Recombinant protein)

[Protocol Download](#)

- ELISA

Gene Info — HLA-DMB

Entrez GeneID

[3109](#)

GeneBank Accession#

[NM_002118](#)

Protein Accession#

[NP_002109](#)

Gene Name

HLA-DMB

Gene Alias

D6S221E, RING7

Gene Description

major histocompatibility complex, class II, DM beta

Omim ID

[142856](#)

Gene Ontology

[Hyperlink](#)

Gene Summary

HLA-DMB belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DMA) and a beta (DMB) chain, both anchored in the membrane. It is located in intracellular vesicles. DM plays a central role in the peptide loading of MHC class II molecules by helping to release the CLIP (class II-associated invariant chain peptide) molecule from the peptide binding site. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. [provided by RefSeq]

Other Designations

MHC class II HLA-DMB|MHC class II antigen HLA-DM beta chain|OTTHUMP00000029257|class II histocompatibility antigen, M beta chain

Pathway

- [Allograft rejection](#)
- [Antigen processing and presentation](#)
- [Asthma](#)
- [Autoimmune thyroid disease](#)
- [Cell adhesion molecules \(CAMs\)](#)
- [Graft-versus-host disease](#)
- [Systemic lupus erythematosus](#)
- [Type I diabetes mellitus](#)

Disease

- [Antiphospholipid Syndrome](#)
- [Arthritis](#)
- [Cardiovascular Diseases](#)
- [Condylomata Acuminata](#)
- [Dermatitis](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)

- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Lupus Erythematosus](#)
- [Measles](#)
- [Psoriasis](#)