


MaxPab®

HNRNPA1 purified MaxPab rabbit polyclonal antibody (D01P)

Catalog # H00003178-D01P Size 100 ug

Applications



Western Blot (Transfected lysate)

Western Blot analysis of HNRNPA1 expression in transfected 293T cell line (<u>H00003178-T05</u>) by HNRNPA1 MaxPab polyclonal antibody.

Lane 1: HNRNPA1 transfected lysate(34.20 KDa).

Lane 2: Non-transfected lysate.

Specification	
Product Description	Rabbit polyclonal antibody raised against a full-length human HNRNPA1 protein.
Immunogen	HNRNPA1 (NP_002127.1, 1 a.a. ~ 320 a.a) full-length human protein.
Sequence	MSKSESPKEPEQLRKLFIGGLSFETTDESLRSHFEQWGTLTDCVVMRDPNTKRSRGFGFVTYAT VEEVDAAMNARPHKVDGRVVEPKRAVSREDSQRPGAHLTVKKIFVGGIKEDTEEHHLRDYFEQY GKIEVIEIMTDRGSGKKRGFAFVTFDDHDSVDKIVIQKYHTVNGHNCEVRKALSKQEMASASSSQR GRSGSGNFGGGRGGGFGGNDNFGRGGNFSGRGGFGGSRGGGGYGGSGDGYNGFGNDGSNFGGGGSYNDFGNYNNQSSNFGPMKGGNFGGRSSGPYGGGGQYFAKPRNQGGYGGSSSSSSYGS GRRF
Host	Rabbit
Reactivity	Human
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.



Applications

Western Blot (Transfected lysate)

Western Blot analysis of HNRNPA1 expression in transfected 293T cell line (<u>H00003178-T05</u>) by HNRNPA1 MaxPab polyclonal antibody.

Lane 1: HNRNPA1 transfected lysate(34.20 KDa).

Lane 2: Non-transfected lysate.

Protocol Download

Entrez GeneID	<u>3178</u>
GeneBank Accession#	NM_002136.2
Protein Accession#	NP_002127.1
Gene Name	HNRNPA1
Gene Alias	HNRPA1, MGC102835
Gene Description	heterogeneous nuclear ribonucleoprotein A1
Omim ID	<u>164017</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene belongs to the A/B subfamily of ubiquitously expressed heterogeneous nuclear ribonucl eoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneo us nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has two repeats of quasi-RRM domains that bind to RNAs. It is one of the most abundant core proteins of hnRNP complexes and it is localized to the nucleoplasm. This protein, along with other hnRNP proteins, is exported from the nucleus, probably bound to mRNA, and is immediately re-imported. Its M9 domain acts as both a nuclear localization and nuclear export signal. The encoded protein is involved in the packaging of pre-mRNA into hnRNP particles, transport of poly A+ mRNA from the nucleus to the cytoplasm, and may modulate splice site selection. It is als o thought have a primary role in the formation of specific myometrial protein species in parturition. Multiple alternatively spliced transcript variants have been found for this gene but only two transcripts are fully described. These variants have multiple alternative transcription initiation sites and multiple polyA sites. [provided by RefSeq
Other Designations	helix-destabilizing protein heterogeneous nuclear ribonucleoprotein A1B protein heterogeneous n

ear ribonucleoprotein particle A1 protein|single-strand DNA-bind

uclear ribonucleoprotein B2 protein|heterogeneous nuclear ribonucleoprotein core protein A1|nucl



Disease

- Cardiovascular Diseases
- Diabetes Mellitus
- Edema