


HOXC10 monoclonal antibody (M02), clone 3F10


Catalog # H00003226-M02 Size 100 ug

Applications



Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged HOXC10 is approximately 0.1ng/ml as a capture antibody.



Western Blot detection against Immunogen (36.63 KDa).

Specification	
Product Description	Mouse monoclonal antibody raised against a partial recombinant HOXC10.
Immunogen	HOXC10 (AAH01293, 158 a.a. ~ 257 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Sequence	LDKTPHCSGANDFEAPFEQRASLNPRAEHLESPQLGGKVSFPETPKSDSQTPSPNEIKTEQSLA GPKGSPSESEKERAKAADSSPDTSDNEAKEEIKAEN
Host	Mouse
Reactivity	Human

😵 Abnova

Product Information

Interspecies Antigen Sequence	Mouse (95)
lsotype	lgG2b Kappa
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Recombinant protein)
 <u>Protocol Download</u>
- Sandwich ELISA (Recombinant protein)
 Detection limit for recombinant GST tagged HOXC10 is approximately 0.1ng/ml as a capture antibody.
 <u>Protocol Download</u>
- ELISA

Gene Info — HOXC10

Entrez GenelD	3226
GeneBank Accession#	<u>BC001293</u>
Protein Accession#	AAH01293
Gene Name	HOXC10
Gene Alias	HOX3I, MGC5259
Gene Description	homeobox C10
Omim ID	605560
Gene Ontology	Hyperlink



Product Information

Gene Summary

This gene belongs to the homeobox family of genes. The homeobox genes encode a highly conse rved family of transcription factors that play an important role in morphogenesis in all multicellular organisms. Mammals possess four similar homeobox gene clusters, HOXA, HOXB, HOXC and H OXD, which are located on different chromosomes and consist of 9 to 11 genes arranged in tand em. This gene is one of several homeobox HOXC genes located in a cluster on chromosome 12. The protein level is controlled during cell differentiation and proliferation, which may indicate this p rotein has a role in origin activation. [provided by RefSeq

Other Designations

homeo box C10|homeobox protein Hox-C10|homeoprotein C10