


IDH2 polyclonal antibody (A01)

Catalog # H00003418-A01


Size 50 uL

Applications



Western Blot (Tissue lysate)

IDH2 polyclonal antibody (A01). Western Blot analysis of IDH2 expression in human ovarian cancer.



Western Blot detection against Immunogen (36.89 kDa) .

Specification

Product Description	Mouse polyclonal antibody raised against a partial recombinant IDH2.
Immunogen	IDH2 (NP_002159, 354 a.a. ~ 451 a.a) partial recombinant protein with GST tag.
Sequence	HYREHQKGRPTSTNPIASIFAWTRGLEHRGKLDGNQDLIRFAQMLEKVCVETVESGAMTKDLAGCI HGLSNVKLNEHFLNTTDFLDTIKSNLDRALGR
Host	Mouse
Reactivity	Human
Interspecies Antigen Sequence	Mouse (97); Rat (97)

Specificity	This antibody cross-reacts with human IDH1.
Quality Control Testing	Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.89 KDa) .
Storage Buffer	50 % glycerol
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Western Blot (Tissue lysate)

IDH2 polyclonal antibody (A01). Western Blot analysis of IDH2 expression in human ovarian cancer.

[Protocol Download](#)

- Western Blot (Recombinant protein)

[Protocol Download](#)

- ELISA

Gene Info — IDH2

Entrez GeneID	3418
GeneBank Accession#	NM_002168
Protein Accession#	NP_002159
Gene Name	IDH2
Gene Alias	ICD-M, IDH, IDHM, IDP, IDPM, mNADP-IDH
Gene Description	isocitrate dehydrogenase 2 (NADP+), mitochondrial
Omim ID	147650
Gene Ontology	Hyperlink

Gene Summary

Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the mitochondria. It plays a role in intermediary metabolism and energy production. This protein may tightly associate or interact with the pyruvate dehydrogenase complex. [provided by RefSeq]

Other Designations

NADP+-specific ICDH|isocitrate dehydrogenase, mitochondrial|oxalosuccinate decarboxylase

Pathway

- [Biosynthesis of alkaloids derived from histidine and purine](#)
- [Biosynthesis of alkaloids derived from ornithine](#)
- [Biosynthesis of alkaloids derived from shikimate pathway](#)
- [Biosynthesis of alkaloids derived from terpenoid and polyketide](#)
- [Biosynthesis of phenylpropanoids](#)
- [Biosynthesis of plant hormones](#)
- [Biosynthesis of terpenoids and steroids](#)
- [Citrate cycle \(TCA cycle\)](#)
- [Glutathione metabolism](#)
- [Metabolic pathways](#)
- [Reductive carboxylate cycle \(CO2 fixation\)](#)

Disease

- [Astrocytoma](#)
- [Blast Crisis](#)
- [Brain Neoplasms](#)
- [Chronic Disease](#)
- [Disease Progression](#)

- [Glioma](#)
- [Hematologic Diseases](#)
- [Leukemia](#)
- [Lung Neoplasms](#)
- [Melanoma](#)
- [Myelodysplastic Syndromes](#)
- [Myeloproliferative Disorders](#)
- [Neoplasm Metastasis](#)
- [Oligodendroglioma](#)
- [Polycythemia Vera](#)
- [Primary Myelofibrosis](#)
- [Recurrence](#)
- [Skin Neoplasms](#)
- [Thrombocythemia](#)