

IFNGR2 rabbit monoclonal antibody

Catalog # H00003460-K

Size 100 ug x up to 3

Specification

Product Description	Rabbit monoclonal antibody raised against a human IFNGR2 peptide using ARM Technology.
Immunogen	A synthetic peptide of human IFNGR2 is used for rabbit immunization. Customer or Abnova will decide on the preferred peptide sequence.
Host	Rabbit
Library Construction	Non-fusion antibody library from rabbit spleen (ARM Technology).
Expression	Overexpression vector and transfection into 293H cell line.
Reactivity	Human
Purification	Protein A
Isotype	IgG
Quality Control Testing	Antibody reactive against human IFNGR2 peptide by ELISA and mammalian transfected lysate by Western Blot.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Deliverable	Up to three rabbit IgG clones of 100 ug each will be delivered to customer.
Note	1. Customer may provide cell or tissue lysate for antibody screening. 2. Rabbit monoclonal antibody generated by ARM technology is amenable to antibody engineering including F(ab) ₂ , IgG, scFv and different Fc and non-Fc conjugates per customer request.

Applications

- Western Blot (Transfected lysate)

[Protocol Download](#)

- ELISA

Gene Info — IFNGR2

Entrez GeneID	3460
GeneBank Accession#	IFNGR2
Gene Name	IFNGR2
Gene Alias	AF-1, IFGR2, IFNGT1
Gene Description	interferon gamma receptor 2 (interferon gamma transducer 1)
Omim ID	147569 209950
Gene Ontology	Hyperlink
Gene Summary	This gene (IFNGR2) encodes the non-ligand-binding beta chain of the gamma interferon receptor. Human interferon-gamma receptor is a heterodimer of IFNGR1 and IFNGR2. Defects in IFNGR2 are a cause of mendelian susceptibility to mycobacterial disease (MSMD), also known as familial disseminated atypical mycobacterial infection. MSMD is a genetically heterogeneous disease with autosomal recessive, autosomal dominant or X-linked inheritance. [provided by RefSeq]
Other Designations	interferon gamma receptor 2 interferon gamma receptor accessory factor-1 interferon gamma receptor beta chain

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Jak-STAT signaling pathway](#)
- [Natural killer cell mediated cytotoxicity](#)

Disease

- [Acute Disease](#)
- [Adenocarcinoma](#)
- [Asthma](#)
- [Atherosclerosis](#)

- [Bronchial Hyperreactivity](#)
- [Bronchiolitis](#)
- [Calcinosis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Chorioamnionitis](#)
- [Coronary Artery Disease](#)
- [Coronary Restenosis](#)
- [Coronary Stenosis](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Drug Hypersensitivity](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Graft vs Host Disease](#)
- [Hepatitis B](#)
- [Hepatitis C](#)
- [Hypersensitivity](#)
- [Infant](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Liver Cirrhosis](#)
- [Lung Neoplasms](#)

- [Lupus Erythematosus](#)
- [Lymphoma](#)
- [Multiple Sclerosis](#)
- [Neovascularization](#)
- [Obstetric Labor](#)
- [Papillomavirus Infections](#)
- [Postoperative Complications](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)
- [Pulmonary Disease](#)
- [Respiratory Syncytial Virus Infections](#)
- [Sarcoidosis](#)
- [Seminoma](#)
- [Stomach Neoplasms](#)
- [Subacute Sclerosing Panencephalitis](#)
- [Testicular Neoplasms](#)
- [Tuberculosis](#)
- [Urinary Bladder Neoplasms](#)
- [Uterine Cervical Neoplasms](#)
- [Viremia](#)
- [Werner syndrome](#)